

# Robust Control and Model Misspecification

Lars Peter Hansen      Thomas J. Sargent  
Gauhar A. Turmuhambetova      Noah Williams\*

September 26, 2005

## Abstract

A decision maker fears that data are generated by a statistical perturbation of an approximating model that is either a controlled diffusion or a controlled measure over continuous functions of time. A perturbation is constrained in terms of its relative entropy. Several different two-player zero-sum games that yield robust decision rules and are related to one another, to the max-min expected utility theory of Gilboa and Schmeidler (1989), and to the recursive risk-sensitivity criterion described in discrete time by Hansen and Sargent (1995). To represent perturbed models, we use martingales on the probability space associated with the approximating model. Alternative sequential and non-sequential versions of robust control theory imply identical robust decision rules that are dynamically consistent in a useful sense.

KEY WORDS: Model uncertainty, entropy, robustness, risk-sensitivity, commitment, time inconsistency, martingale.

## 1 Introduction

A *decision maker* consists of (i) a utility function that is maximized subject to (ii) a model. Classical decision and control theory assume that a decision maker has complete confidence in his model. Robust control theory presents alternative formulations of a decision maker who doubts his model. To capture the idea that the decision maker views his model as an approximation, these formulations alter items (i) and (ii) by (1) surrounding the decision maker's approximating model with a cloud of models that are difficult to distinguish with finite data, and (2) adding a malevolent second agent. The malevolent agent promotes

---

\*We thank Fernando Alvarez, David Backus, Gary Chamberlain, Ivar Ekeland, Peter Klibanoff, Tomasz Piskorski, Michael Allen Rierson, Aldo Rustichini, Jose Scheinkman, Christopher Sims, Nizar Touzi, and especially Costis Skiadas for valuable comments on an earlier draft. Sherwin Rosen encouraged us to write this paper.

robustness by causing the decision maker to explore the fragility of candidate decision rules to departures of the data from the approximating model. Finding a rule that is robust to model misspecification entails computing lower bounds on a rule's performance. The minimizing agent constructs those lower bounds.

Different parts of robust control theory uses alternative mathematical formalisms. While all of them have versions of items (1) and (2), they differ in many important mathematical details including the probability spaces on which they are defined; their ways of representing alternative models; their restrictions on sets of alternative models; and their protocols about the timing of choices by the maximizing and minimizing decision makers. Nevertheless, common outcomes and representations emerge from all of these alternative formulations. Equivalent concerns about model misspecification can be represented by either (a) altering the decision maker's preferences to enhance risk-sensitivity, or (b) leaving his preferences alone but slanting his expectations relative to his approximating model in a particular context-specific way, or (c) adding a set of perturbed models and a malevolent agent. This paper exhibits these unifying connections and stresses how they can be exploited in applications.

Robust control theory shares with both the Bayesian paradigm and the rational expectations model the feature that the decision maker brings to the table one fully specified model. In robust control theory it is called either his reference model or his approximating model. Although the decision maker does not explicitly specify alternative models, he evaluates a decision rule under a set of incompletely articulated models that are formed by perturbing his approximating model. Robust control theory contributes thoughtful ways to surround a single approximating model with a cloud of other models. We give technical conditions that allow us to regard that set of models as the multiple priors that appear in the max-min expected utility theory of Gilboa and Schmeidler (1989). Some technical conditions allow us to represent the approximating model and perturbations to it. Other technical conditions reconcile the equilibrium outcomes of several two-player zero-sum games that have different timing protocols, providing a way of interpreting robust control in terms of a recursive version of max-min expected utility theory.

This paper starts with two alternative ways of representing an approximating model in continuous time – either (1) as a diffusion or (2) as a measure over continuous functions of time that are induced by the diffusion. We consider different ways of perturbing each such representation of the approximating model. These lead to alternative formulations of robust control problems. In all of our problems, we use a definition of relative entropy (an expected log likelihood ratio) to constrain the gap between the approximating model and a statistical perturbation to it. We take the maximum value of that gap as a parameter that measures the set of perturbations against which the decision maker seeks robustness. Requiring that entropy be finite restricts the form that model misspecification can take. In particular, finiteness of entropy implies that admissible perturbations of the approximating model must be absolutely continuous with respect to it over finite intervals. For a diffusion, absolute continuity over finite intervals implies that allowable perturbations can alter the drift but not the volatility. Restricting ourselves to perturbations that are absolutely continuous over finite intervals is therefore tantamount to considering perturbed models that are in principle

statistically difficult to distinguish from the approximating model, an idea exploited by Anderson, Hansen, and Sargent (2003) to calibrate a plausible amount of fear of model misspecification in a study of market prices of risk.

The work of Araujo and Sandroni (1999) and Sandroni (2000) emphasizes that absolute continuity of models implies that decision makers' beliefs eventually merge with the model that generates the data. But in infinite horizon economies, absolute continuity over finite intervals does not imply absolute continuity. By allowing perturbations that are not absolutely continuous, we arrest the merging of models and thereby create a setting in which a decision maker's fear of model misspecification endures. Perturbations that are absolutely continuous over finite intervals but still not absolutely continuous can be difficult to detect from a continuous record of finite length, though they could be detected from a continuous data record of infinite length. We discuss how this modeling choice interacts with the way that the decision maker discounts the future.

We also consider a variety of technical issues about timing protocols that underlie interconnections among various expressions of robust control theory. A Bellman-Isaacs condition allows us to exchange orders of minimization and maximization and validates several useful results, including the existence of a Bayesian interpretation of a robust decision rule.

Counterparts to many of the issues treated in this paper occur in discrete time robust control theory. Many of these issues surface in nonstochastic versions of the theory, for example, in Başar and Bernhard (1995). The continuous time stochastic setting of this paper allows sharper analytical results in several cases.

## 1.1 Language

We call a problem *nonsequential* if, at an initial time 0, a decision maker chooses an entire history-contingent sequence. We call a problem *sequential* or *recursive* if, at each time  $t \geq 0$ , a decision maker chooses the time  $t$  component of his action process as a function of his time  $t$  information.

## 1.2 Organization of paper

The technical nature of interrelated material inspires us to present it in two exposures consisting first of section 2, then of the remaining sections. Section 2 sets aside a variety of complications and compiles our main results by displaying Hamilton-Jacobi-Bellman (HJB) equations for various games and decision problems and asserting without proof the key relationships among them. The remaining sections lay things out in detail. Section 3 sets the stage by describing both sequential and nonsequential versions of an ordinary control problem under a known model. These problems form benchmarks against which to judge subsequent problems in which the decision maker distrusts his model. Section 3 also introduces a risk-sensitive control problem that alters the decision maker's objective function but leaves unchallenged his trust in his model. Section 4 discusses alternative ways of representing fear of model misspecification. Section 5 introduces entropy and its relationship

to a concept of absolute continuity over finite intervals, then formulates two nonsequential zero-sum two-player games, called penalty and constraint games, that induce robust decision rules. The games in section 5 are both cast in terms of sets of probability measures. In section 6, we cast counterparts to these games on a fixed probability measure by representing perturbations to an approximating model in terms of martingales defined on a fixed probability space. Section 7 gives a sequential formulation of a penalty game. By taking continuation entropy as an endogenous state variable, section 8 gives a sequential formulation of a constraint game. This formulation sets the stage for our discussion in section 9 of the dynamic consistency issues raised by Epstein and Schneider (2004). Section 10 concludes. Appendix A presents the cast of characters that records the objects and concepts that occur throughout the paper. Four additional appendixes deliver proofs.

## 2 Overview

One Hamilton-Jacobi-Bellman (HJB) equation is worth a thousand words. This section concisely summarizes our main results by displaying HJB equations for various two-player zero-sum continuous time games that are defined in terms of a Markov diffusion with state  $x$  and Brownian motion  $B$ , together with the value functions for some related nonsequential games. Our story is encoded in state variables, drifts, and diffusion terms that occur in HJB equations for several optimum problems and dynamic games. This telegraphic section is intended for readers who glean everything from HJB equations and as a summary of key findings. Readers who prefer a more deliberate presentation from the beginning should skip to section 3.

### 2.1 Sequential control problems and games

#### Benchmark control problem:

We take as a benchmark an ordinary control problem with value function

$$J(x_0) = \max_{c \in C} E \left[ \int_0^\infty \exp(-\delta t) U(c_t, x_t) dt \right]$$

where the maximization is subject to  $dx_t = \mu(c_t, x_t)dt + \sigma(c_t, x_t)dB_t$  and where  $x_0$  is a given initial condition. The HJB equation for the benchmark problem is

$$\delta J(\tilde{x}) = \max_{\tilde{c} \in \tilde{C}} U(\tilde{c}, \tilde{x}) + \mu(\tilde{c}, \tilde{x}) \cdot J_x(\tilde{x}) + \frac{1}{2} \text{trace} [\sigma(\tilde{c}, \tilde{x})' J_{xx}(\tilde{x}) \sigma(\tilde{c}, \tilde{x})]. \quad (1)$$

The notation  $\tilde{\cdot}$  is used to denote a potentially realized value of a control or a state. Similarly,  $\tilde{C}$  is the set of admissible values for the control. Subscripts on value functions denote the respective derivatives. We provide more detail about the benchmark problem in section 3.1.

In the benchmark problem, the decision maker trusts his model. We want to study comparable problems where the decision maker distrusts his model. Several superficially different

devices can be used to promote robustness to misspecification of the diffusion associated with (1). These add either a free parameter  $\theta > 0$  or a state variable  $\tilde{r} \geq 0$  or a state vector  $X$  and produce recursive problems with one of the following HJB equations:

**Risk sensitive control problem:**

$$\begin{aligned} \delta S(\tilde{x}) = & \max_{\check{c} \in \check{C}} U(\check{c}, \tilde{x}) + \mu(\check{c}, \tilde{x}) \cdot S_x(\tilde{x}) + \frac{1}{2} \text{trace} [\sigma(\check{c}, \tilde{x})' S_{xx}(\tilde{x}) \sigma(\check{c}, \tilde{x})] \\ & - \frac{1}{2\theta} S_x(\tilde{x})' \sigma(\check{c}, \tilde{x}) \sigma(\check{c}, \tilde{x})' S_x(\tilde{x}) \end{aligned} \quad (2)$$

HJB equation (2) alters the right side of the value function recursion (1) by deducting  $\frac{1}{2\theta}$  times the local variation of the continuation value. The optimal decision rule for the risk-sensitive problem (2) is a policy function

$$c_t = \alpha_c(x_t)$$

where the dependence on  $\theta$  is understood. In control theory,  $-1/\theta$  is called the *risk-sensitivity parameter*; in the recursive utility literature, it is called the *variance multiplier*. Section 3.2 below provides more details about the risk-sensitive problem.

**Penalty robust control problem:**

A two-player zero-sum game has a value function  $M$  that satisfies

$$M(\tilde{x}, \tilde{z}) = \tilde{z}V(\tilde{x})$$

where  $z_t$  is another state variable that changes the probability distribution and  $V$  satisfies the HJB equation:

$$\begin{aligned} \delta V(\tilde{x}) = & \max_{\check{c} \in \check{C}} \min_{\check{h}} U(\check{c}, \tilde{x}) + \frac{\theta}{2} \check{h} \cdot \check{h} + [\mu(\check{c}, \tilde{x}) + \sigma(\check{c}, \tilde{x})\check{h}] \cdot V_x(\tilde{x}) \\ & + \frac{1}{2} \text{trace} [\sigma(\check{c}, \tilde{x})' V_{xx}(\tilde{x}) \sigma(\check{c}, \tilde{x})]. \end{aligned} \quad (3)$$

The process  $z = \{z_t : t \geq 0\}$  is a martingale with initial condition  $z_0 = 1$  and evolution  $dz_t = h_t \cdot dB_t$ . The minimizing agent in (3) chooses an  $\check{h}$  to alter the probability distribution;  $\theta > 0$  is a parameter that penalizes the minimizing agent for distorting the drift. Optimizing over  $\check{h}$  shows that  $V$  from (3) solves the same partial differential equation (2). The penalty robust control problem is discussed in more detail in sections 6.4 and 7.

**Constraint robust control problem:**

A two-player zero-sum game has a value function  $\tilde{z}K(\tilde{x}, \tilde{r})$ , where  $K$  satisfies the HJB equation

$$\begin{aligned} \delta K(\check{x}, \check{r}) &= \max_{\check{c} \in \check{C}} \min_{\check{h}, \check{g}} U(\check{c}, \check{x}) + [\mu(\check{c}, \check{x}) + \sigma(\check{c}, \check{x})\check{h}] \cdot K_x(\check{x}, \check{r}) + \left( \delta \check{r} - \frac{\check{h} \cdot \check{h}}{2} \right) \cdot K_r(\check{x}, \check{r}) \\ &\quad + \frac{1}{2} \text{trace} \left( \begin{bmatrix} \sigma(\check{c}, \check{x})' & \check{g} \end{bmatrix} \begin{bmatrix} K_{xx}(\check{x}, \check{r}) & K_{xr}(\check{x}, \check{r}) \\ K_{rx}(\check{x}, \check{r}) & K_{rr}(\check{x}, \check{r}) \end{bmatrix} \begin{bmatrix} \sigma(\check{c}, \check{x}) \\ \check{g}' \end{bmatrix} \right). \end{aligned} \quad (4)$$

Equation (4) shares with (3) that the minimizing agent chooses an  $\check{h}$  that alters the probability distribution, but unlike (3), there is no penalty parameter  $\theta$ . Instead, in (4), the minimizing agent's choice of  $h_t$  affects a new state variable  $r_t$  that we call *continuation entropy*. The minimizing player also controls another decision variable  $\check{g}$  that determines how increments in the continuation value are related to the underlying Brownian motion. The right side of the HJB equation for the constraint control problem (4) is attained by decision rules

$$c_t = \phi_c(x_t, r_t), \quad h_t = \phi_h(x_t, r_t), \quad g_t = \phi_g(x_t, r_t).$$

We can solve the equation  $\frac{\partial}{\partial r} K(x_t, r_t) = -\theta$  to express  $r_t$  as a time invariant function of  $x_t$ :

$$r_t = \phi_r(x_t).$$

Therefore, along an equilibrium path of game (4), we have  $c_t = \phi_c[x_t, \phi_r(x_t)]$ ,  $h_t = \phi_h[x_t, \phi_r(x_t)]$ ,  $g_t = \phi_g[x_t, \phi_r(x_t)]$ . More detail on the constraint problem is given in section 8.

### A problem with a Bayesian interpretation:

A single agent optimization problem has a value function  $\check{z}W(\check{x}, \check{X})$  where  $W$  satisfies the HJB equation:

$$\begin{aligned} \delta W(\check{x}, \check{X}) &= \max_{\check{c} \in \check{C}} U(\check{c}, \check{x}) + \mu(\check{c}, \check{x}) \cdot W_x(\check{x}, \check{X}) + \mu^*(\check{x}) \cdot W_X(\check{x}, \check{X}) \\ &\quad + \frac{1}{2} \text{trace} \left( \begin{bmatrix} \sigma(\check{c}, \check{x})' & \sigma^*(\check{X})' \end{bmatrix} \begin{bmatrix} W_{xx}(\check{x}, \check{X}) & W_{xX}(\check{x}, \check{X}) \\ W_{Xx}(\check{x}, \check{X}) & W_{XX}(\check{x}, \check{X}) \end{bmatrix} \begin{bmatrix} \sigma(\check{c}, \check{x}) \\ \sigma^*(\check{X}) \end{bmatrix} \right) \\ &\quad + \alpha_h(\check{X}) \cdot \sigma(\check{c}, \check{x})' W_x(\check{x}, \check{X}) + \alpha_h(\check{X}) \cdot \sigma^*(\check{X})' W_X(\check{x}, \check{X}) \end{aligned} \quad (5)$$

where  $\mu^*(\check{X}) = \mu[\alpha_c(\check{X}), \check{X}]$  and  $\sigma^*(\check{X}) = \sigma[\alpha_c(\check{X}), \check{X}]$ . The function  $W(\check{x}, \check{X})$  in (5) depends on an additional component of the state vector  $\check{X}$  that is comparable in dimension with  $\check{x}$  and that is to be initialized from the common value  $\check{X}_0 = \check{x}_0 = x_0$ . We shall show in appendix E that equation (5) is the HJB equation for an ordinary (i.e., single agent) control problem with discounted objective:

$$z_0 W(\check{x}, \check{X}) = E \int_0^\infty \exp(-\delta t) z_t U(c_t, x_t) dt$$

and state evolution:

$$\begin{aligned} dx_t &= \mu(c_t, x_t) dt + \sigma(c_t, x_t) dB_t \\ dz_t &= z_t \alpha_h(X_t) dB_t \\ dX_t &= \mu^*(X_t) dt + \sigma^*(X_t) dB_t \end{aligned}$$

with  $z_0 = 1$ ,  $x_0 = \check{x}$ , and  $X_0 = \check{X}$ .

This problem alters the benchmark control problem by changing the probabilities assigned to the *shock* process  $\{B_t : t \geq 0\}$ . It differs from the penalty robust control problem (3) because the process  $z$  used to change probabilities does not depend on state variables that are endogenous to the control problem.

In appendix E, we verify that under the optimal  $c$  and the prescribed choices of  $\mu^*$ ,  $\sigma^*$ ,  $\alpha_h$ , the ‘big X’ component of the state vector equals the ‘little x’ component, provided that  $\check{X}_0 = \check{x}_0$ . Equation (5) is therefore the HJB equation for an ordinary control problem that justifies a robust decision rule under a fixed probability model that differs from the approximating model. As the presence of  $z_t$  as a preference shock suggests, this problem reinterprets the equilibrium of the two-player zero-sum game portrayed in the penalty robust control problem (3). For a given  $\theta$  that gets embedded in  $\sigma^*$ ,  $\mu^*$ , the right side of the HJB equation (5) is attained by  $\check{c} = \gamma_c(\check{x}, \check{X})$ .

## 2.2 Different ways to attain robustness

Relative to (1), HJB equations (2), (3), (4), and (5) can all be interpreted as devices that in different ways promote robustness to misspecification of the diffusion. HJB equations (2) and (5) are for ordinary control problems: only the maximization operator appears on the right side, so that there is no minimizing player to promote robustness. Problem (2) promotes robustness by enhancing the maximizing player’s sensitivity to risk, while problem (5) promotes robustness by attributing to the maximizing player a belief about the state transition law that is distorted in a pessimistic way relative to his approximating model. The HJB equations in (3) and (4) describe two-player zero-sum dynamic games in which a minimizing player promotes robustness.

## 2.3 Nonsequential problems

We also study two nonsequential two-player zero-sum games that are defined in terms of perturbations  $q \in Q$  to the measure  $q^0$  over continuous functions of time that is induced by the Brownian motion  $B$  in the diffusion for  $x$ . Let  $q_t$  be the restriction of  $q$  to events measurable with respect to time  $t$  histories of observations. We define discounted relative entropy as

$$\tilde{\mathcal{R}}(q) \doteq \delta \int_0^\infty \exp(-\delta t) \left( \int \log \left( \frac{dq_t}{dq_t^0} \right) dq_t \right) dt$$

and use it to restrict the size of perturbations  $q$  to  $q^0$ . Leaving the dependence on  $B$  implicit, we define a utility process  $v_t(c) = U(c_t, x_t)$  and pose the following two problems:

**Nonsequential penalty control problem:**

$$\tilde{V}(\theta) = \max_{c \in \mathcal{C}} \min_{q \in Q} \int_0^\infty \exp(-\delta t) \left( \int v_t(c) dq_t \right) dt + \theta \tilde{\mathcal{R}}(q). \quad (6)$$

**Nonsequential constraint control problem:**

$$\tilde{K}(\eta) = \max_{c \in \mathcal{C}} \min_{q \in Q(\eta)} \int_0^\infty \exp(-\delta t) \left( \int v_t(c) dq_t \right) dt \quad (7)$$

where  $Q(\eta) = \{q \in Q : \tilde{\mathcal{R}}(q) \leq \eta\}$ .

Problem (7) fits the max-min expected utility model of Gilboa and Schmeidler (1989), where  $Q(\eta)$  is a set of multiple priors. The axiomatic treatment of Gilboa and Schmeidler views this set of priors as an expression of the decision maker's preferences and does not cast them as perturbations of an approximating model.<sup>1</sup> We are free to think of problem (7) as providing a way to use a single approximating model  $q^0$  to generate Gilboa-Schmeidler's set of priors as all those unspecified models that satisfy the restriction on relative entropy,  $Q(\eta) = \{q \in Q : \tilde{\mathcal{R}}(q) \leq \eta\}$ . In section 5 we provide more detail on the nonsequential problems.

The objective functions for these two nonsequential optimization problems (6) and (7) are related via the Legendre transform pair:

$$\tilde{V}(\theta) = \min_{\eta \geq 0} \tilde{K}(\eta) + \theta\eta \quad (8)$$

$$\tilde{K}(\eta) = \max_{\theta \geq 0} \tilde{V}(\theta) - \eta\theta. \quad (9)$$

## 2.4 Connections

An association between robust control and the framework of Gilboa and Schmeidler (1989) extends beyond problem (7) because the equilibrium value functions and decision rules for all of our problems are intimately related. Where  $V$  is the value function in (3) and  $K$  is the value function in (4), the recursive counterpart to (8) is:

$$V(\tilde{x}) = \min_{\tilde{r} \geq 0} K(\tilde{x}, \tilde{r}) + \theta\tilde{r}$$

with the implied first-order condition

$$\frac{\partial}{\partial \tilde{r}} K(\tilde{x}, \tilde{r}) = -\theta.$$

This first-order condition implicitly defines  $\tilde{r}$  as a function of  $\tilde{x}$  for a given  $\theta$ , which implies that  $\tilde{r}$  is a redundant state variable. The penalty formulation avoids this redundancy.<sup>2</sup>

The nonsequential value function  $\tilde{V}$  is related to the other value functions via:

$$\tilde{V}(\theta) = M(x_0, 1) = 1 \cdot V(x_0) = W(x_0, x_0) = S(x_0)$$

---

<sup>1</sup>Similarly, Savage's framework does not purport to describe the process by which the Bayesian decision maker constructs his unique prior.

<sup>2</sup>There is also a recursive analog to (9) that uses the fact that the function  $V$  depends implicitly on  $\theta$ .

where  $x_0$  is the common initial value and  $\theta$  is held fixed across the different problems. Though these problems have different decision rules, we shall show that for a fixed  $\theta$  and comparable initial conditions, they have identical equilibrium outcomes and identical recursive representations of those outcomes. In particular, the following relations prevail across the equilibrium decision rules for our different problems:

$$\alpha_c(\tilde{x}) = \gamma_c(\tilde{x}, \tilde{x}) = \phi_c[\tilde{x}, \phi_r(\tilde{x})].$$

## 2.5 Who cares?

We care about the equivalence of these control problems and games because some of the problems are easier to solve and others are easier to interpret.

These problems came from literatures that approached the problem of decision making in the presence of model misspecification from different angles. The recursive version of the penalty problem (3) emerged from a literature on robust control that also considered the risk-sensitive problem (2). The nonsequential constraint problem (7) is an example of the min-max expected utility theory of Gilboa and Schmeidler (1989) with a particular set of priors. By modifying the set of priors over time, constraint problem (4) states a recursive version of that nonsequential constraint problem. The Lagrange multiplier theorem supplies an interpretation of the penalty parameter  $\theta$ .

A potentially troublesome feature of multiple priors models for applied work is that they impute a *set* of models to the decision maker.<sup>3</sup> How should that set be specified? Robust control theory gives a convenient way to specify and measure a set of priors surrounding a single approximating model.

## 3 Three ordinary control problems

By describing three ordinary control problems, this section begins describing the technical conditions that underlie the broad claims made in section 2. In each problem, a single decision maker chooses a stochastic process to maximize an intertemporal return function. The first two are different representations of the same underlying problem. They are cast on different probability spaces and express different timing protocols. The third, called the risk-sensitive control problem, alters the objective function of the decision maker to induce more aversion to risk.

### 3.1 Benchmark problem

We start with two versions of a benchmark stochastic optimal control problem. The first formulation is defined in terms of a state vector  $x$ , an underlying probability space  $(\Omega, \mathcal{F}, P)$ , a  $d$ -dimensional, standard Brownian motion  $\{B_t : t \geq 0\}$  defined on that space, and  $\{\mathcal{F}_t :$

---

<sup>3</sup>For applied work, an attractive feature of rational expectations is that by equating the equilibrium of the model itself to the decision maker's prior, decision makers' beliefs contribute no free parameters.

$t \geq 0\}$ , the completion of the filtration generated by the Brownian motion  $B$ . For any stochastic process  $\{a_t : t \geq 0\}$ , we use  $a$  or  $\{a_t\}$  to denote the process and  $a_t$  to denote the time  $t$ -component of that process. The random vector  $a_t$  maps  $\Omega$  into a set  $\check{A}$ ;  $\check{a}$  denotes an element in  $\check{A}$ . Actions of the decision-maker form a progressively measurable stochastic process  $\{c_t : t \geq 0\}$ , which means that the time  $t$  component  $c_t$  is  $\mathcal{F}_t$  measurable.<sup>4</sup> Let  $U$  be an instantaneous utility function and  $C$  be the set of admissible control processes.

**Definition 3.1.** *The benchmark control problem is:*

$$J(x_0) = \sup_{c \in C} E \left[ \int_0^\infty \exp(-\delta t) U(c_t, x_t) dt \right] \quad (10)$$

where the maximization is subject to

$$dx_t = \mu(c_t, x_t)dt + \sigma(c_t, x_t)dB_t \quad (11)$$

and where  $x_0$  is a given initial condition.

The parameter  $\delta$  is a subjective discount rate,  $\mu$  is the drift coefficient and  $\sigma\sigma'$  is the diffusion matrix. We restrict  $\mu$  and  $\sigma$  so that any progressively measurable control  $c$  in  $C$  implies a progressively measurable state vector process  $x$  and maintain

**Assumption 3.2.**  *$J(x_0)$  is finite.*

We shall refer to the law of motion (11) or the probability measure over sequences that it induces as the decision maker's *approximating model*. The benchmark control problem treats the approximating model as correct.

### 3.1.1 A nonsequential version of the benchmark problem

It is useful to restate the benchmark problem in terms of the probability space that the Brownian motion induces over continuous functions of time, thereby converting it into a nonsequential problem that pushes the state  $x$  into the background. At the same time, it puts the induced probability distribution in the foreground and features the linearity of the objective in the induced probability distribution. For similar constructions and further discussions of induced distributions, see Elliott (1982) and Liptser and Shiryaev (2000), chapter 7.

The  $d$ -dimensional Brownian motion  $B$  induces a multivariate Wiener measure  $q^0$  on a *canonical space*  $(\Omega^*, \mathcal{F}^*)$ , where  $\Omega^*$  is the space of continuous functions  $f : [0, +\infty) \rightarrow \mathbb{R}^d$  and  $\mathcal{F}_t^*$  is the Borel sigma algebra for the restriction of the continuous functions  $f$  to  $[0, t]$ . Define open sets using the sup-norm over each interval. Notice that  $\iota_s(f) \doteq f(s)$  is  $\mathcal{F}_t^*$

---

<sup>4</sup>Progressive measurability requires that we view  $c \doteq \{c_t : t \geq 0\}$  as a function of  $(t, \omega)$ . For any  $t \geq 0$ ,  $c : [0, t] \times \Omega$  must be measurable with respect to  $\mathcal{B}_t \times \mathcal{F}_t$ , where  $\mathcal{B}_t$  is a collection of Borel subsets of  $[0, t]$ . See Karatzas and Shreve (1991) pages 4 and 5 for a discussion.

measurable for each  $0 \leq s \leq t$ . Let  $\mathcal{F}^*$  be the smallest sigma algebra containing  $\mathcal{F}_t^*$  for  $t \geq 0$ . An event in  $\mathcal{F}_t^*$  restricts continuous functions on the finite interval  $[0, t]$ . For any probability measure  $q$  on  $(\Omega^*, \mathcal{F}^*)$ , let  $q_t$  denote the restriction to  $\mathcal{F}_t^*$ . In particular,  $q_t^0$  is the multivariate Wiener measure over the event collection  $\mathcal{F}_t^*$ .

Given a progressively measurable control  $c$ , solve the stochastic differential equation (11) to obtain a progressively measurable utility process

$$U(c_t, x_t) = v_t(c, B)$$

where  $v(c, \cdot)$  is a progressively measurable family defined on  $(\Omega^*, \mathcal{F}^*)$ . This notation accounts for but conceals the evolution of the state vector  $x_t$ . A realization of the Brownian motion is a continuous function. Putting a probability measure  $q^0$  on the space of continuous functions allows us to evaluate expectations. We leave implicit the dependence on  $B$  and represent the decision maker's objective as  $\int_0^\infty \exp(-\delta t) \left( \int v_t(c) dq_t^0 \right) dt$ .

**Definition 3.3.** *A nonsequential benchmark control problem is*

$$\tilde{J}(x_0) = \sup_{c \in C} \int_0^\infty \exp(-\delta t) \left( \int v_t(c) dq_t^0 \right) dt.$$

### 3.1.2 Recursive version of the benchmark problem

The problem in definition 3.1 asks the decision maker once and for all at time 0 to choose an entire process  $c \in C$ . To transform the problem into one in which the decision maker chooses sequentially, we impose additional structure on the choice set  $C$  by restricting  $\check{c}$  to be in some set  $\check{C}$  that is common for all dates. This is for notational simplicity, since we could easily incorporate control constraints of the form  $\check{C}(t, \check{x})$ . With this specification of controls, we make the problem recursive by asking the decision maker to choose  $\check{c}$  as a function of the state  $x$  at each date.

**Definition 3.4.** *The HJB equation for the benchmark problem is*

$$\delta J(\check{x}) = \sup_{\check{c} \in \check{C}} U(\check{c}, \check{x}) + \mu(\check{c}, \check{x}) \cdot J_x(\check{x}) + \frac{1}{2} \text{trace} [\sigma(\check{c}, \check{x})' J_{xx}(\check{x}) \sigma(\check{c}, \check{x})]. \quad (12)$$

The recursive version of the benchmark problem (12) puts the state  $x_t$  front and center. A decision rule  $c_t = \zeta_c(x_t)$  attains the right side of the HJB equation (12).

Although the nonsequential and recursive versions of the benchmark control problem yield identical formulas for  $(c, x)$  as a function of the Brownian motion  $B$ , they differ in how they represent the same approximating model: as a probability distribution in the nonsequential problem as a stochastic differential equation in the recursive problem. Both versions of the benchmark problem treat the decision maker's approximating model as true.<sup>5</sup>

<sup>5</sup>As we discuss more in section 7, an additional argument is generally needed to show that an appropriate solution of (12) is equal to the value of the original problem (10).

### 3.2 Risk-sensitive control

Let  $\rho$  be an intertemporal return or utility function. Instead of maximizing  $E\rho$  (where  $E$  continues to mean mathematical expectation), risk-sensitive control theory maximizes  $-\theta \log E[\exp(-\rho/\theta)]$ , where  $1/\theta$  is a risk-sensitivity parameter. As the name suggests, the exponentiation inside the expectation makes this objective more sensitive to risky outcomes. Jacobson (1973) and Whittle (1981) initiated risk sensitive optimal control in the context of discrete-time linear-quadratic decision problems. Jacobson and Whittle showed that the risk-sensitive control law can be computed by solving a robust penalty problem of the type we have studied here.

A *risk-sensitive* control problem treats the decision maker's approximating model as true but alters preferences by appending an additional term to the right side of the HJB equation (12):

$$\begin{aligned} \delta S(\tilde{x}) &= \sup_{\tilde{c} \in \tilde{C}} U(\tilde{c}, \tilde{x}) + \mu(\tilde{c}, \tilde{x}) \cdot S_x(\tilde{x}) + \frac{1}{2} \text{trace} [\sigma(\tilde{c}, \tilde{x})' S_{xx}(\tilde{x}) \sigma(\tilde{c}, \tilde{x})] \\ &\quad - \frac{1}{2\theta} S_x(\tilde{x})' \sigma(\tilde{c}, \tilde{x}) \sigma(\tilde{c}, \tilde{x})' S_x(\tilde{x}), \end{aligned} \tag{13}$$

where  $\theta > 0$ . The term

$$\mu(\tilde{c}, \tilde{x}) \cdot S_x(\tilde{x}) + \frac{1}{2} \text{trace} [\sigma(\tilde{c}, \tilde{x})' S_{xx}(\tilde{x}) \sigma(\tilde{c}, \tilde{x})]$$

in HJB equation (13) is the local mean or  $dt$  contribution to the continuation value process  $\{S(x_t) : t \geq 0\}$ . Thus, (13) adds  $-\frac{1}{2\theta} S_x(\tilde{x})' \sigma(\tilde{c}, \tilde{x}) \sigma(\tilde{c}, \tilde{x})' S_x(\tilde{x})$  to the right side of the HJB equation for the benchmark control problem (10), (11). Notice that  $S_x(x_t)' \sigma(c_t, x_t) dB_t$  gives the local Brownian contribution to the value function process  $\{S(x_t) : t \geq 0\}$ . The additional term in the HJB equation is the negative of the local variance of the continuation value weighted by  $\frac{1}{2\theta}$ . Relative to our discussion above, we can view this as the Ito's lemma correction term for the evolution of instantaneous expected utility that comes from the concavity of the exponentiation in the risk sensitive objective. When  $\theta = +\infty$ , this collapses to the benchmark control problem. When  $\theta < \infty$ , we call it a risk-sensitive control problem with  $-\frac{1}{\theta}$  being the *risk-sensitivity* parameter. A solution of the risk-sensitive control problem is attained by a policy function

$$c_t = \alpha_c(x_t) \tag{14}$$

whose dependence on  $\theta$  is understood.

James (1992) studied a continuous-time, nonlinear diffusion formulation of a risk-sensitive control problem. Risk-sensitive control theory typically focuses on the case in which the discount rate  $\delta$  is zero. Hansen and Sargent (1995) showed how to introduce discounting and still preserve much of the mathematical structure for the linear-quadratic, Gaussian risk-sensitive control problem. They applied the recursive utility framework developed by Epstein and Zin (1989) in which the risk-sensitive adjustment is applied recursively to the continuation values. Recursive formulation (13) gives the continuous-time counterpart for Markov diffusion processes. Duffie and Epstein (1992) characterized the preferences that underlie this specification.

## 4 Fear of model misspecification

For a given  $\theta$ , the optimal risk-sensitive decision rule emerges from other problems in which the decision maker’s objective function remains that in the benchmark problem (10) and in which the adjustment to the continuation value in (13) reflects not altered preferences but distrust of the model (11). Moreover, just as we formulated the benchmark problem either as a nonsequential problem with induced distributions or as a recursive problem, there are also nonsequential and recursive representations of robust control problems.

Each of our decision problems for promoting robustness to model misspecification is a zero-sum, two-player game in which a maximizing player (‘the decision maker’) chooses a best response to a malevolent player (‘nature’) who can alter the stochastic process within prescribed limits. The minimizing player’s malevolence is the maximizing player’s tool for analyzing the fragility of alternative decision rules. Each game uses a Nash equilibrium concept. We portray games that differ from one another in three dimensions: (1) the protocols that govern the timing of players’ decisions, (2) the constraints on the malevolent player’s choice of models; and (3) the mathematical spaces in terms of which the games are posed. Because the state spaces and probability spaces on which they are defined differ, the recursive versions of these problems yield decision rules that differ from (14). Despite that, all of the formulations give rise to identical decision processes for  $c$ , all of which in turn are equal to those that apply the optimal risk sensitive decision rule (14) to the transition equation (11).

The equivalence of their outcomes provides interesting alternative perspectives from which to understand the decision maker’s response to possible model misspecification.<sup>6</sup> That outcomes are identical for these different games means that when all is said and done, the timing protocols don’t matter. Because some of the timing protocols correspond to nonsequential or ‘static’ games while others enable sequential choices, equivalence of equilibrium outcomes implies a form of dynamic consistency.

Jacobson (1973) and Whittle (1981) first showed that the risk-sensitive control law can be computed by solving a robust penalty problem of the type we have studied here, but without discounting. Subsequent research reconfirmed this link in nonsequential and undiscounted problems, typically posed in nonstochastic environments. Petersen, James, and Dupuis (2000) explicitly considered an environment with randomness, but did not make the link to recursive risk-sensitivity.

## 5 Two robust control problems defined on sets of probability measures

We formalize the connection between two problems that are robust counterparts to the nonsequential version of the benchmark control problem (3.3). These problems do not fix an induced probability distribution  $q^o$ . Instead they express alternative models as alternative

---

<sup>6</sup>See section 9 of Anderson, Hansen, and Sargent (2003) for an application.

induced probability distributions and add a player who chooses a probability distribution to minimize the objective. This leads to a pair of two-player zero-sum games. One of the two games falls naturally into the framework of Gilboa and Schmeidler (1989) and the other is closely linked to risk-sensitive control. An advantage of working with the induced distributions is that a convexity property that helps to establish the connection between the two games is easy to demonstrate.

## 5.1 Entropy and absolute continuity over finite intervals

We use a notion of absolute continuity of one infinite-time stochastic process with respect to another that is weaker than what is implied by the standard definition of absolute continuity. The standard notion characterizes two stochastic processes as being absolutely continuous with respect to each other if they agree about “tail events”. Roughly speaking, the weaker concept requires that the two measures being compared both put positive probability on all of the same events, *except* tail events. This weaker notion of absolute continuity is interesting for applied work because of what it implies about how quickly it is possible statistically to distinguish one model from another.

Recall that the Brownian motion  $B$  induces a multivariate Wiener measure on  $(\Omega^*, \mathcal{F}^*)$  that we have denoted  $q^0$ . For any probability measure  $q$  on  $(\Omega^*, \mathcal{F}^*)$ , we have let  $q_t$  denote the restriction to  $\mathcal{F}_t^*$ . In particular,  $q_t^0$  is the multivariate Wiener measure over the events  $\mathcal{F}_t^*$ .

**Definition 5.1.** *A distribution  $q$  is said to be absolutely continuous over finite intervals with respect to  $q^0$  if  $q_t$  is absolutely continuous with respect to  $q_t^0$  for all  $t < \infty$ .*<sup>7</sup>

Let  $Q$  be the set of all distributions that are absolutely continuous with respect to  $q^0$  over finite intervals. The set  $Q$  is convex. Absolute continuity over finite intervals captures the idea that two models are difficult to distinguish given samples of finite length. If  $q$  is absolutely continuous with respect to  $q^0$  over finite intervals, we can construct likelihood ratios for finite histories at any calendar date  $t$ . To measure the discrepancy between models over an infinite horizon, we use a discounted measure of *relative entropy*:

$$\tilde{\mathcal{R}}(q) \doteq \delta \int_0^\infty \exp(-\delta t) \left( \int \log \left( \frac{dq_t}{dq_t^0} \right) dq_t \right) dt, \quad (15)$$

where  $\frac{dq_t}{dq_t^0}$  is the Radon-Nikodym derivative of  $q_t$  with respect to  $q_t^0$ . In appendix B (claim B.1), we show that this discrepancy measure is convex in  $q$ .

The distribution  $q$  is absolutely continuous with respect to  $q^0$  when

$$\int \log \left( \frac{dq}{dq^0} \right) dq < +\infty.$$

---

<sup>7</sup>Kabanov, Lipcer, and Sirjaev (1979) refer to this concept as *local* absolute continuity. Although Kabanov, Lipcer, and Sirjaev (1979) define local absolute continuity through the use of stopping times, they argue that their definition is equivalent to this “simpler one”.

In this case a law of large numbers that applies under  $q_0$  must also apply under  $q$ , so that discrepancies between them are at most ‘temporary’. We introduce discounting in part to provide an alternative interpretation of the recursive formulation of risk-sensitive control as expressing a fear of model misspecification rather than extra aversion to well understood risks. By restricting the discounted entropy (15) to be finite, we allow

$$\int \log \left( \frac{dq}{dq^0} \right) dq = +\infty. \quad (16)$$

Time series averages of functions that converge almost surely under  $q^0$  can converge to a different limit under  $q$ , or they may not converge at all. That would allow a statistician to distinguish  $q$  from  $q^0$  with a continuous record of data on an infinite interval.<sup>8</sup> But we want these alternative models to be close enough to the approximating model that they are statistically difficult to distinguish from it after having observed a continuous data record of only finite length  $N$  on the state. We implement this requirement by requiring  $\tilde{\mathcal{R}}(q) < +\infty$ , where  $\tilde{\mathcal{R}}(q)$  is defined in (15).

The presence of discounting in (15) and its absence from (16) are significant. With alternative models that satisfy (16), the decision maker seeks robustness against models that can be distinguished from the approximating model with an infinite data record; but because the models satisfy (15), it is difficult to distinguish them from a finite data record. Thus, we have in mind settings of  $\delta$  for which impatience outweighs the decision maker’s ability eventually to learn specifications that give superior fits, prompting him to focus on designing a robust decision rule.

We now have the vocabulary to state two nonsequential robust control problems that use  $Q$  as a family of distortions to the probability distribution  $q^0$  in the benchmark problem:

**Definition 5.2.** A nonsequential penalty robust control problem is

$$\tilde{V}(\theta) = \sup_{c \in C} \inf_{q \in Q} \int_0^\infty \exp(-\delta t) \left( \int v_t(c) dq_t \right) dt + \theta \tilde{\mathcal{R}}(q).$$

**Definition 5.3.** A nonsequential constraint robust control problem is

$$\tilde{K}(\eta) = \sup_{c \in C} \inf_{q \in Q(\eta)} \int_0^\infty \exp(-\delta t) \left( \int v_t(c) dq_t \right) dt$$

where  $Q(\eta) = \{q \in Q : \tilde{\mathcal{R}}(q) \leq \eta\}$ .

The first problem is closely linked to the risk sensitive control problem. The second problem fits into the max-min expected utility or multiple priors model advocated by Gilboa and Schmeidler (1989), the set of priors being  $Q(\eta)$ . We use  $\theta$  to index a family of penalty robust control problems and  $\eta$  to index a family of constraint robust control problems. The two types of problems are linked by the Lagrange multiplier theorem, as we show next.

---

<sup>8</sup>Our specification allows  $Q$  measures to put different probabilities on tail events, which prevents the conditional measures from merging, as Blackwell and Dubins (1962) show will occur under absolute continuity. See Kalai and Lerner (1993) and Jackson, Kalai, and Smorodinsky (1999) for implications of absolute continuity for learning.

## 5.2 Relation between the constraint and penalty problems

In this subsection we establish two important things about the two nonsequential multiple priors problems 5.2 and 5.3: (1) we show that we can interpret the robustness parameter  $\theta$  in problem 5.2 as a Lagrange multiplier on the specification-error constraint  $\tilde{\mathcal{R}}(q) \leq \eta$  in problem 5.3;<sup>9</sup> (2) we display technical conditions that make the solutions of the two problems equivalent to one another. We shall exploit both of these results in later sections.

The simultaneous maximization and minimization means that the link between the penalty and constraint problem is not a direct implication of the Lagrange multiplier Theorem. The following treatment exploits convexity of  $\tilde{R}$  in  $Q$ . The analysis follows Petersen, James, and Dupuis (2000), although our measure of entropy differs.<sup>10</sup> As in Petersen, James, and Dupuis (2000), we use tools of convex analysis contained in Luenberger (1969) to establish the connection between the two problems.

Assumption 3.2 makes the optimized objectives for both the penalty and constraint robust control problems less than  $+\infty$ . They can be  $-\infty$ , depending on the magnitudes of  $\theta$  and  $\eta$ .

Given an  $\eta^* > 0$ , add  $-\theta\eta^*$  to the objective in problem 5.2. For given  $\theta$ , doing this has no impact on the control law.<sup>11</sup> For a given  $c$ , the objective of the constraint robust control problem is linear in  $q$  and the entropy measure  $\tilde{\mathcal{R}}$  in the constraint is convex in  $q$ . Moreover, the family of admissible probability distributions  $Q$  is itself convex. Thus, we formulate the constraint version of the robust control problem (problem 5.3) as a Lagrangian:

$$\sup_{c \in C} \inf_{q \in Q} \sup_{\theta \geq 0} \int_0^\infty \exp(-\delta t) \left( \int v_t(c) dq_t \right) dt + \theta \left[ \tilde{\mathcal{R}}(q) - \eta \right].$$

For many choices of  $q$ , The optimizing multiplier  $\theta$  is degenerate: it is infinite if  $q$  violates the constraint and zero if the constraint is slack. Therefore, we include  $\theta = +\infty$  in the choice set for  $\theta$ . Exchanging the order of  $\max_\theta$  and  $\min_q$  attains the same value of  $q$ . The Lagrange multiplier theorem allows us to study:

$$\sup_{c \in C} \sup_{\theta \geq 0} \inf_{q \in Q} \int_0^\infty \exp(-\delta t) \left( \int v_t(c) dq_t \right) dt + \theta \left[ \tilde{\mathcal{R}}(q) - \eta \right]. \quad (17)$$

A complication arises at this point because the maximizing  $\theta$  in (17) depends on the choice of  $c$ . In solving a robust control problem, we are most interested in the  $c$  that solves the constraint robust control problem. We can find the appropriate choice of  $\theta$  by changing the order of  $\max_c$  and  $\max_\theta$  to obtain:

$$\sup_{\theta \geq 0} \sup_{c \in C} \inf_{q \in Q} \int_0^\infty \exp(-\delta t) \left( \int v_t(c) dq_t \right) dt + \theta \left[ \tilde{\mathcal{R}}(q) - \eta^* \right] = \max_{\theta \geq 0} \tilde{V}(\theta) - \theta\eta^*,$$

<sup>9</sup>This connection is regarded as self-evident throughout the literature on robust control. It has been explored in the context of a linear-quadratic control problem, informally by Hansen, Sargent, and Tallarini (1999), and formally by Hansen and Sargent (2006).

<sup>10</sup>To accommodate discounting in the recursive, risk sensitive control problem, we include discounting in our measure of entropy. See appendix B.

<sup>11</sup>However, it will alter which  $\theta$  results in the highest objective.

since for a given  $\theta$  the term  $-\theta\eta^*$  does not effect the extremizing choices of  $(c, q)$ .

**Claim 5.4.** *For  $\eta^* > 0$ , suppose that  $c^*$  and  $q^*$  solve the constraint robust control problem for  $\tilde{K}(\eta^*) > -\infty$ . Then there exists a  $\theta^* > 0$  such that the corresponding penalty robust control problem has the same solution. Moreover,*

$$\tilde{K}(\eta^*) = \max_{\theta \geq 0} \tilde{V}(\theta) - \theta\eta^*.$$

*Proof.* This result is essentially the same as Theorem 2.1 of Petersen, James, and Dupuis (2000) and follows directly from Luenberger (1969).  $\square$

This claim gives  $\tilde{K}$  as the Legendre transform of  $\tilde{V}$ . Moreover, by adapting an argument of Luenberger (1969), we can show that  $\tilde{K}$  is decreasing and convex in  $\eta$ .<sup>12</sup> We are interested in recovering  $\tilde{V}$  from  $\tilde{K}$  as the inverse Legendre transform via:

$$\tilde{V}(\theta^*) = \min_{\eta \geq 0} \tilde{K}(\eta) + \theta^*\eta. \quad (18)$$

It remains to justify this recovery formula.

We call *admissible* those nonnegative values of  $\theta$  for which it is feasible to make the objective function greater than  $-\infty$ . If  $\hat{\theta}$  is admissible, values of  $\theta$  larger than  $\hat{\theta}$  are also admissible, since these values only make the objective larger. Let  $\underline{\theta}$  denote the greatest lower bound for admissible values of  $\theta$ . Consider a value  $\theta^* > \underline{\theta}$ . Our aim is to find a constraint associated with this choice of  $\theta$ .

It follows from claim 5.4 that

$$\tilde{V}(\theta^*) \leq \tilde{K}(\eta) + \theta^*\eta$$

for any  $\eta > 0$  and hence

$$\tilde{V}(\theta^*) \leq \min_{\eta \geq 0} \tilde{K}(\eta) + \theta^*\eta.$$

Moreover,

$$\tilde{K}(\eta) \leq \inf_{q \in Q(\eta)} \sup_{c \in C} \int_0^\infty \exp(-\delta t) \left( \int v_t(c) dq_t \right) dt,$$

since maximizing *after* minimizing (rather than vice versa) cannot decrease the resulting value of the objective. Thus,

$$\begin{aligned} \tilde{V}(\theta^*) &\leq \min_{\eta \geq 0} \left[ \inf_{q \in Q(\eta)} \sup_{c \in C} \int_0^\infty \exp(-\delta t) \left( \int v_t(c) dq_t \right) dt + \theta^*\eta \right] \\ &= \min_{\eta \geq 0} \left[ \inf_{q \in Q(\eta)} \sup_{c \in C} \int_0^\infty \exp(-\delta t) \left( \int v_t(c) dq_t \right) dt + \theta^*\tilde{\mathcal{R}}(q) \right] \\ &= \inf_{q \in Q} \sup_{c \in C} \int_0^\infty \exp(-\delta t) \left( \int v_t(c) dq_t \right) dt + \theta^*\tilde{\mathcal{R}}(q). \end{aligned}$$

---

<sup>12</sup>This follows because we may view  $\tilde{K}$  as the maximum over convex functions indexed by alternative consumption processes.

For the first equality, the minimization over  $\eta$  is important. Given some  $\hat{\eta}$  we may lower the objective by substituting  $\tilde{\mathcal{R}}(q)$  for  $\hat{\eta}$  when the constraint  $\tilde{\mathcal{R}}(q) \leq \hat{\eta}$  is imposed in the inner minimization problem. Thus the minimized choice of  $q$  for  $\hat{\eta}$  may have entropy  $\tilde{\eta} < \hat{\eta}$ . More generally, there may exist a sequence  $\{q_j : j = 1, 2, \dots\}$  that approximates the inf for which  $\{\tilde{\mathcal{R}}(q_j) : j = 1, 2, \dots\}$  is bounded away from  $\hat{\eta}$ . In this case we may extract a subsequence of  $\{\mathcal{R}(q_j) : j = 1, 2, \dots\}$  that converges to  $\tilde{\eta} < \hat{\eta}$ . Therefore, we would obtain the same objective by imposing an entropy constraint  $\tilde{\mathcal{R}}(q) \leq \tilde{\eta}$  at the outset:

$$\begin{aligned} & \inf_{q \in Q(\tilde{\eta})} \left[ \sup_{c \in C} \int_0^\infty \exp(-\delta t) \left( \int v_t(c) dq_t \right) dt + \theta^* \tilde{\eta} \right] \\ &= \inf_{q \in Q(\tilde{\eta})} \left[ \sup_{c \in C} \int_0^\infty \exp(-\delta t) \left( \int v_t(c) dq_t \right) dt + \theta^* \tilde{\mathcal{R}}(q) \right]. \end{aligned}$$

Since the objective is minimized by choice  $\eta$  there is no further reduction in the optimized objective by substituting  $\tilde{\mathcal{R}}(q)$  for  $\eta$ .

Notice that the last equality gives a min – max analogue to the *nonsequential penalty problem* (5.2), but with the order of minimization and maximization reversed. If the resulting value continues to be  $\tilde{V}(\theta^*)$ , we have verified (18).

We shall invoke the following assumption:

**Assumption 5.5.** For  $\theta > \underline{\theta}$

$$\begin{aligned} \tilde{V}(\theta) &= \max_{c \in C} \min_{q \in Q} \int_0^\infty \exp(-\delta t) \left( \int v_t(c) dq_t \right) dt + \theta \tilde{\mathcal{R}}(q) \\ &= \min_{q \in Q} \max_{c \in C} \int_0^\infty \exp(-\delta t) \left( \int v_t(c) dq_t \right) dt + \theta \tilde{\mathcal{R}}(q). \end{aligned}$$

Both equalities assume that the maximum and minimum are attained. Because minimization occurs first, without the assumption the second equality would have to be replaced by a less than or equal sign ( $\leq$ ). In much of what follows, we presume that inf's and sup's are attained in the control problems, and thus we will replace inf with min and sup with max.

**Claim 5.6.** Suppose that Assumption 5.5 is satisfied and that for  $\theta^* > \underline{\theta}$ ,  $c^*$  is the maximizing choice of  $c$  for the penalty robust control problem 5.2. Then that  $c^*$  also solves the constraint robust control problem 5.3 for  $\eta^* = \tilde{\mathcal{R}}(q^*)$  where  $\eta^*$  solves

$$\tilde{V}(\theta^*) = \min_{\eta \geq 0} \tilde{K}(\eta) + \theta^* \eta.$$

Since  $\tilde{K}$  is decreasing and convex,  $\tilde{V}$  is increasing and concave in  $\theta$ . The Legendre and inverse Legendre transforms given in claims 5.4 and 5.6 fully describe the mapping between the constraint index  $\eta^*$  and the penalty parameter  $\theta^*$ . However, given  $\eta^*$ , they do not imply that the associated  $\theta^*$  is unique, nor for a given  $\theta^* > \underline{\theta}$  do they imply that the associated  $\eta^*$  is unique.

While claim 5.6 maintains assumption 5.5, claim 5.4 does not. Without assumption 5.5, we do not have a proof that  $\tilde{V}$  is concave. Moreover, for some values of  $\theta^*$  and a solution pair

$(c^*, q^*)$  of the penalty problem, we may not be able to produce a corresponding constraint problem. Nevertheless, the family of penalty problems indexed by  $\theta$  continues to embed the solutions to the constraint problems indexed by  $\eta$  as justified by claim 5.4. We are primarily interested in problems for which assumption 5.5 is satisfied and in section 7 and appendix D provide some sufficient conditions for this assumption. One reason for interest in this assumption is given in the next subsection.

### 5.3 Preference Orderings

We now define two preference orderings associated with the constraint and penalty control problems. One preference ordering uses the value function:

$$\hat{K}(c; \eta) = \inf_{\tilde{\mathcal{R}}(q) \leq \eta} \int_0^\infty \exp(-\delta t) \left( \int v_t(c) dq_t \right) dt.$$

**Definition 5.7.** (*Constraint preference ordering*) For any two progressively measurable  $c$  and  $c^*$ ,  $c^* \succeq_\eta c$  if

$$\hat{K}(c^*; \eta) \geq \hat{K}(c; \eta).$$

The other preference ordering uses the value function:

$$\hat{V}(c; \theta) = \inf_q \int_0^\infty \exp(-\delta t) \left( \int v_t(c) dq_t \right) dt + \theta \tilde{\mathcal{R}}(q)$$

**Definition 5.8.** (*Penalty preference ordering*) For any two progressively measurable  $c$  and  $c^*$ ,  $c^* \succeq_\theta c$  if

$$\tilde{V}(c^*; \theta) \geq \tilde{V}(c; \theta).$$

The first preference order has the multiple-priors form justified by Gilboa and Schmeidler (1989). The second is commonly used to compute robust decision rules and is closest to recursive utility theory. The two preference orderings differ. Furthermore, given  $\eta$ , there exists no  $\theta$  that makes the two preference orderings agree. However, the Lagrange Multiplier Theorem delivers a weaker result that is very useful to us. While they differ globally, indifference curves passing through a given point  $c^*$  in the consumption set are tangent for the two preference orderings. For asset pricing, a particularly interesting point  $c^*$  would be one that solves an optimal resource allocation problem.

Use the Lagrange Multiplier Theorem to write  $\hat{K}$  as

$$\hat{K}(c^*; \eta^*) = \max_{\theta \geq 0} \inf_q \int_0^\infty \exp(-\delta t) \left( \int v_t(c^*) dq_t \right) dt + \theta \left[ \tilde{\mathcal{R}}(q) - \eta^* \right],$$

and let  $\theta^*$  denote the maximizing value of  $\theta$ , which we assume to be strictly positive. Suppose that  $c^* \succeq_{\eta^*} c$ . Then

$$\hat{V}(c; \theta^*) - \theta^* \eta^* \leq \hat{K}(c; \eta^*) \leq \hat{K}(c^*; \eta^*) = \hat{V}(c^*; \theta^*) - \theta^* \eta^*.$$

Thus,  $c^* \succeq_{\theta^*} c$ . The observational equivalence results from Claims 5.4 and 5.6 apply to decision profile  $c^*$ . The indifference curves touch but do not cross at this point.

Although the preferences differ, the penalty preferences are of interest in their own right. See Wang (2001) for an axiomatic development of entropy-based preference orders and Maccheroni, Marinacci, and Rustichini (2004) for an axiomatic treatment of preferences specified using convex penalization.

## 5.4 Bayesian interpretation of outcome of nonsequential game

A widespread device for interpreting a statistical decision rule is to find a probability distribution for which the decision rule is optimal. Here we seek an induced probability distribution for  $B$  such that the solution for  $c$  from either the constraint or penalty robust decision problem is optimal for a counterpart to the benchmark problem. When we can produce such a distribution, we say that we have a Bayesian interpretation for the robust decision rule. (See Blackwell and Girshick (1954) and Chamberlain (2000) for related discussions.)

The freedom to exchange orders of maximization and minimization in problem 5.2 (Assumption 5.5) justifies such a Bayesian interpretation of the decision process  $c \in C$ . Let  $(c^*, q^*)$  be the equilibrium of game 5.2. Given the worst case model  $q^*$ , consider the control problem:

$$\max_{c \in C} \int_0^\infty \exp(-\delta t) \left( \int v_t(c) dq_t^* \right) dt. \quad (19)$$

Problem (19) is a version of our nonsequential benchmark problem 3.3 with a fixed model  $q^*$  that is distorted relative to the approximating model  $q^0$ . The optimal choice of a progressively measurable  $c$  takes  $q^*$  as exogenous. The optimal decision  $c^*$  is not altered by adding  $\theta \tilde{\mathcal{R}}(q^*)$  to the objective. Therefore, being able to exchange orders of extremization in 5.2 allows us to support a solution to the penalty problem by a particular distortion in the Wiener measure. The implied least favorable  $q^*$  assigns a different (induced) probability measure for the exogenous stochastic process  $\{B_t : t \geq 0\}$ . Given that distribution,  $c^*$  is the ordinary (non robust) optimal control process.

Having connected the penalty and the constraint problem, in what follows we will focus primarily on the penalty problem. For notational simplicity, we will simply fix a value of  $\theta$  and not formally index a family of problems by this parameter value.

## 6 Games on fixed probability spaces

This section describes important technical details that are involved in moving from the nonsequential to the recursive versions of the multiple probability games 5.2 and 5.3. It is convenient to represent alternative model specifications as martingale ‘preference shocks’ on a common probability space. This allows us to formulate two-player zero-sum differential games and to use existing results for such games. Thus, instead of working with multiple

distributions on the measurable space  $(\Omega^*, \mathcal{F}^*)$ , we now use the original probability space  $(\Omega, \mathcal{F}, P)$  in conjunction with nonnegative martingales.

We present a convenient way to parameterize the martingales and issue a caveat about this parameterization.

## 6.1 Martingales and finite interval absolute continuity

For any continuous function  $f$  in  $\Omega^*$ , let

$$\begin{aligned}\kappa_t(f) &= \left( \frac{dq_t}{dq_t^0} \right) (f) \\ z_t &= \kappa_t(B)\end{aligned}\tag{20}$$

where  $\kappa_t$  is the Radon-Nikodym derivative of  $q_t$  with respect to  $q_t^0$ .

**Claim 6.1.** *Suppose that for all  $t \geq 0$ ,  $q_t$  is absolutely continuous with respect to  $q_t^0$ . The process  $\{z_t : t \geq 0\}$  defined via (20) on  $(\Omega, \mathcal{F}, P)$  is a nonnegative martingale adapted to the filtration  $\{\mathcal{F}_t : t \geq 0\}$  with  $Ez_t = 1$ . Moreover,*

$$\int \phi_t dq_t = E[z_t \phi_t(B)]\tag{21}$$

for any bounded and  $\mathcal{F}_t^*$  measurable function  $\phi_t$ . Conversely, if  $\{z_t : t \geq 0\}$  is a nonnegative progressively measurable martingale with  $Ez_t = 1$ , then the probability measure  $q$  defined via (21) is absolutely continuous with respect to  $q^0$  over finite intervals.

*Proof.* The first part of this claim follows directly from the proof of theorem 7.5 in Liptser and Shiryaev (2000). Their proof is essentially a direct application of the Law of Iterated Expectations and the fact that probability distributions necessarily integrate to one. Conversely, suppose that  $z$  is a nonnegative martingale on  $(\Omega, \mathcal{F}, P)$  with unit expectation. Let  $\phi_t$  be any nonnegative, bounded and  $\mathcal{F}_t^*$  measurable function. Then (21) defines a measure because indicator functions are nonnegative, bounded functions. Clearly  $\int \phi_t dq_t = 0$  whenever  $E\phi_t(B) = 0$ . Thus,  $q_t$  is absolutely continuous with respect to  $q_t^0$ , the measure induced by Brownian motion restricted to  $[0, t]$ . Setting  $\phi_t = 1$  shows that  $q_t$  is in fact a probability measure for any  $t$ .  $\square$

Claim 6.1 is important because it allows us to integrate over  $(\Omega^*, \mathcal{F}^*, q)$  by instead integrating against a martingale  $z$  on the original probability space  $(\Omega, \mathcal{F}, P)$ .

## 6.2 Representing martingales

By exploiting the Brownian motion information structure, we can attain a convenient representation of a martingale. Any martingale  $z$  with a unit expectation can be portrayed as

$$z_t = 1 + \int_0^t k_u dB_u$$

where  $k$  is a progressively measurable  $d$ -dimensional process that satisfies:

$$P \left\{ \int_0^t |k_u|^2 du < \infty \right\} = 1$$

for any finite  $t$  (see Revuz and Yor (1994), Theorem V.3.4). Define:

$$h_t = \begin{cases} k_t/z_t & \text{if } z_t > 0 \\ 0 & \text{if } z_t = 0. \end{cases} \quad (22)$$

Then  $z$  solves the integral equation

$$z_t = 1 + \int_0^t z_u h_u dB_u \quad (23)$$

and its differential counterpart

$$dz_t = z_t h_t dB_t \quad (24)$$

with initial condition  $z_0 = 1$ , where for  $t > 0$

$$P \left\{ \int_0^t (z_u)^2 |h_u|^2 du < \infty \right\} = 1. \quad (25)$$

The scaling by  $(z_u)^2$  permits

$$\int_0^t |h_u|^2 du = \infty$$

provided that  $z_t = 0$  on the probability one event in (25).

In reformulating the nonsequential penalty problem 5.2, we parameterize nonnegative martingales by progressively measurable processes  $h$ . We introduce a new state  $z_t$  initialized at one, and take  $h$  to be under the control of the minimizing agent.

## 6.3 Representing likelihood ratios

We are now equipped to fill in some important details associated with using martingales to represent likelihood ratios for dynamic models. Before addressing these issues, we use a simple static example to exhibit an important idea.

### 6.3.1 A static example

The static example is designed to illustrate two alternative ways to represent the expected value of a likelihood ratio by changing the measure with respect to which it is evaluated. Consider two models of a vector  $y$ . In the first,  $y$  is normally distributed with mean  $\nu$  and covariance matrix  $I$ . In the second,  $y$  is normally distributed with mean zero and covariance matrix  $I$ . The logarithm of the ratio of the first density to the second is:

$$\ell(y) = \left( \nu \cdot y - \frac{1}{2} \nu \cdot \nu \right).$$

Let  $E^1$  denote the expectation under model one and  $E^2$  under model two. Properties of the log-normal distribution imply that

$$E^1 \exp[\ell(y)] = 1.$$

Under the second model

$$E^2 \ell(y) = E^1 \ell(y) \exp[\ell(y)] = \frac{1}{2} \nu \cdot \nu,$$

which is relative entropy.

### 6.3.2 The dynamic counterpart

We now consider a dynamic counterpart to the static example by showing two ways to represent likelihood ratios, one under the original Brownian motion model and another under the model associated with a nonnegative martingale  $z$ . First we consider the likelihood ratio under the Brownian motion model for  $B$ . As noted above, the solution to (24) can be represented as an exponential:

$$z_t = \exp \left( \int_0^t h_u \cdot dB_u - \frac{1}{2} \int_0^t |h_u|^2 du \right). \quad (26)$$

We allow  $\int_0^t |h_u|^2 du$  to be infinite with positive probability and adopt the convention that the exponential is zero when this event happens. In the event that  $\int_0^t |h_u|^2 du < \infty$ , we can define the stochastic integral  $\int_0^t h_u dB_u$  as an appropriate probability limit (see Lemma 6.2 of Liptser and Shiryaev (2000)).

When  $z$  is a martingale, we can interpret the right side of (26) as a formula for the likelihood ratio of two models evaluated under the Brownian motion specification for  $B$ . Taking logarithms, we find that

$$\ell_t = \int_0^t h_u \cdot dB_u - \frac{1}{2} \int_0^t |h_u|^2 du.$$

Since  $h$  is progressively measurable, we can write:

$$h_t = \psi_t(B).$$

Changing the distribution of  $B$  in accordance with  $q$  gives another characterization of the likelihood ratio. The Girsanov Theorem implies

**Claim 6.2.** *If for all  $t \geq 0$ ,  $q_t$  is absolutely continuous with respect to  $q_t^0$ , then  $q$  is the induced distribution for a (possibly weak) solution  $B$  to a stochastic differential equation defined on a probability space  $(\Omega, \mathcal{F}, \tilde{P})$ :*

$$dB_t = \psi_t(B)dt + d\tilde{B}_t$$

for some progressively measurable  $\psi$  defined on  $(\Omega^*, \mathcal{F}^*)$  and some Brownian motion  $\tilde{B}$  that is adapted to  $\{\mathcal{F}_t : t \geq 0\}$ . Moreover, for each  $t$

$$\tilde{P} \left[ \int_0^t |\psi_u(B)|^2 du < \infty \right] = 1.$$

*Proof.* From Lemma 6.1 there is a nonnegative martingale  $z$  associated with the Radon-Nikodym derivative of  $q_t$  with respect to  $q_t^0$ . This martingale has expectation unity for all  $t$ . The conclusion follows from a generalization of the Girsanov Theorem (*e.g.* see Liptser and Shiryaev (2000) Theorem 6.2).  $\square$

The  $\psi_t(B)$  is the same as that used to represent  $h_t$  defined by (22). Under the distribution  $\tilde{P}$ ,

$$B_t = \int_0^t h_u du + \tilde{B}_t$$

where  $\tilde{B}_t$  is a Brownian motion with respect to the filtration  $\{\mathcal{F}_t : t \geq 0\}$ . In other words, we obtain perturbed models by replacing the Brownian motion model for a shock process with a Brownian motion with a drift.

Using this representation, we can write the logarithm of the likelihood ratio as:

$$\tilde{\ell}_t = \int_0^t \psi_u(B) \cdot d\tilde{B}_u + \frac{1}{2} \int_0^t |\psi_u(B)|^2 du.$$

**Claim 6.3.** For  $q \in Q$ , let  $z$  be the nonnegative martingale associated with  $q$  and let  $h$  be the progressively measurable process satisfying (23). Then

$$\tilde{\mathcal{R}}(q) = \frac{1}{2} E \left[ \int_0^\infty \exp(-\delta t) z_t |h_t|^2 dt \right].$$

*Proof.* See appendix B.  $\square$

This claim leads us to define a discounted entropy measure for nonnegative martingales:

$$\mathcal{R}^*(z) \doteq \frac{1}{2} E \left[ \int_0^\infty \exp(-\delta t) z_t |h_t|^2 dt \right]. \quad (27)$$

## 6.4 A martingale version of a robust control problem

Modeling alternative probability distributions as preference shocks that are martingales on a common probability space is mathematically convenient because it allows us to reformulate the penalty robust control problem (problem 5.2) as:

**Definition 6.4.** A nonsequential martingale robust control problem is

$$\max_{c \in \mathcal{C}} \min_{h \in H} E \left( \int_0^\infty \exp(-\delta t) z_t \left[ U(c_t, x_t) + \frac{\theta}{2} |h_t|^2 \right] dt \right) \quad (28)$$

subject to:

$$\begin{aligned} dx_t &= \mu(c_t, x_t)dt + \sigma(c_t, x_t)dB_t \\ dz_t &= z_t h_t \cdot dB_t. \end{aligned} \tag{29}$$

But there is potentially a technical problem with this formulation. There may exist control process  $h$  and corresponding processes  $z$  such that  $z$  is a nonnegative local martingale for which  $\mathcal{R}^*(z) < \infty$ , yet  $z$  is not a martingale. We have not ruled out nonnegative supermartingales that happen to be local martingales. This means that even though  $z$  is a local martingale, it might satisfy only the inequality

$$E(z_t | \mathcal{F}_s) \leq z_s$$

for  $0 < s \leq t$ . Even when we initialize  $z_0$  to one,  $z_t$  may have a mean less than one and the corresponding measure will not be a probability measure. Then we would have given the minimizing agent more options than we intend.

For this not to cause difficulty, at the very least we have to show that the minimizing player's choice of  $h$  in problem 6.4 is associated with a  $z$  that is a martingale and not just a supermartingale.<sup>13</sup> More generally, we have to verify that enlarging the set of processes  $z$  as we have done does not alter the equilibrium of the two-player zero-sum game. In particular, consider the second problem in assumption 5.5. It suffices to show that the minimizing  $h$  implies a  $z$  that is a martingale. If we assume that condition 5.5 is satisfied, then it suffices to check this for the following timing protocol:

$$\min_{h \in H} \max_{c \in C} E \left( \int_0^\infty \exp(-\delta t) z_t \left[ U(c_t, x_t) + \frac{\theta}{2} |h_t|^2 \right] dt \right)$$

subject to (29),  $z_0 = 1$ , and an initial condition  $x_0$  for  $x$ .<sup>14</sup> In appendix C, we show how to establish that the solution is indeed a martingale.

---

<sup>13</sup>Alternatively, we might interpret the supermartingale as allowing for an escape to a terminal absorbing state with a terminal value function equal to zero. The expectation of  $z_t$  gives the probability that an escape has not happened as of date  $t$ . The existence of such terminal state is not, however, entertained in our formulation of 5.2.

<sup>14</sup>To see this let  $H^* \subseteq H$  be the set of controls  $h$  for which  $z$  is a martingale and let  $\text{obj}(h, c)$  be the objective as a function of the controls. Then under Assumption 5.5 we have

$$\min_{h \in H^*} \max_{c \in C} \text{obj}(h, c) \geq \min_{h \in H} \max_{c \in C} \text{obj}(h, c) = \max_{c \in C} \min_{h \in H} \text{obj}(h, c) \leq \max_{c \in C} \min_{h \in H^*} \text{obj}(h, c). \tag{30}$$

If we demonstrate, the first inequality  $\geq$  in (30) is an equality, it follows that

$$\min_{h \in H^*} \max_{c \in C} \text{obj}(h, c) \leq \max_{c \in C} \min_{h \in H^*} \text{obj}(h, c).$$

Since the reverse inequality is always satisfied provided that the extrema are attained, this inequality can be replaced by an equality. It follows that the second inequality  $\leq$  in (30) must in fact be an equality as well.

## 7 Sequential timing protocol for a penalty formulation

The martingale problem 6.4 assumes that at time zero both decision makers commit to decision processes whose time  $t$  components are measurable functions of  $\mathcal{F}_t$ . The minimizing decision maker who chooses distorted beliefs  $h$  takes  $c$  as given; and the maximizing decision maker who chooses  $c$  takes  $h$  as given. Assumption 5.5 asserts that the order in which the two decision makers choose does not matter.

This section studies a two-player zero-sum game with a protocol that makes both players choose sequentially. We set forth conditions that imply that with sequential choices we obtain the same time zero value function and the same outcome path that would prevail were both players to choose once and for all at time 0. The sequential formulation is convenient computationally and also gives a way to justify the exchange of orders of extremization stipulated by assumption 5.5.

We have used  $c$  to denote the control process and  $\check{c} \in \check{C}$  to denote the value of a control at a particular date. We let  $\check{h} \in \check{H}$  denote the realized martingale control at any particular date. We can think of  $\check{h}$  as a vector in  $\mathbb{R}^d$ . Similarly, we think of  $\check{x}$  and  $\check{z}$  as being realized states.

To analyze outcomes under a sequential timing protocol, we think of varying the initial state and define a value function  $M(x_0, z_0)$  as the optimized objective function (28) for the martingale problem. By appealing to results of Fleming and Souganidis (1989), we can verify that  $\tilde{V}(\theta) = M(\check{x}, \check{z}) = \check{z}V(\check{x})$ , provided that  $\check{x} = x_0$  and  $\check{z} = 1$ . Under a sequential timing protocol, this same value function gives the *continuation value* for evaluating states reached at subsequent time periods.

Fleming and Souganidis (1989) show that a *Bellman-Isaacs condition* renders equilibrium outcomes under two-sided commitment at date zero identical with outcomes of a Markov perfect equilibrium in which the decision rules of both agents are chosen sequentially, each as a function of the state vector  $x_t$ .<sup>15</sup> The HJB equation for the infinite-horizon zero-sum two-player martingale game is:

$$\begin{aligned} \delta \check{z}V(\check{x}) &= \max_{\check{c} \in \check{C}} \min_h \check{z}U(\check{c}, \check{x}) + \check{z} \frac{\theta}{2} \check{h} \cdot \check{h} + \mu(\check{c}, \check{x}) \cdot V_x(\check{x}) \check{z} \\ &\quad + \check{z} \frac{1}{2} \text{trace} [\sigma(\check{c}, \check{x})' V_{xx}(\check{x}) \sigma(\check{c}, \check{x})] + \check{z} \check{h} \cdot \sigma(\check{c}, \check{x})' V_x(\check{x}) \end{aligned} \quad (31)$$

where  $V_x$  is the vector of partial derivatives of  $V$  with respect to  $\check{x}$  and  $V_{xx}$  is the matrix of second derivatives.<sup>16</sup> The diffusion specification makes this HJB equation a partial differential equation that has multiple solutions that correspond to different boundary conditions.

<sup>15</sup>Fleming and Souganidis (1989) impose as restrictions that  $\mu$ ,  $\sigma$  and  $U$  are bounded, uniformly continuous and Lipschitz continuous with respect to  $\check{x}$  uniformly in  $\check{c}$ . They also require that the controls  $\check{c}$  and  $\check{h}$  reside in compact sets. While these restrictions are imposed to obtain general existence results, they are not satisfied for some important examples. Presumably existence in these examples will require special arguments. These issues are beyond the scope of this paper.

<sup>16</sup>In general the value functions associated with stochastic control problems will not be twice differentiable,

To find the true value function and to justify the associated control laws requires that we apply a *Verification Theorem* (e.g. see Theorem 5.1 of Fleming and Soner (1993)).

The scaling of partial differential equation (31) by  $\check{z}$  verifies our guess that the value function is linear in  $z$ . This allows us to study the alternative HJB equation:

$$\begin{aligned} \delta V(\check{x}) &= \max_{\check{c} \in \check{C}} \min_{\check{h}} U(\check{c}, \check{x}) + \frac{\theta}{2} \check{h} \cdot \check{h} + [\mu(\check{c}, \check{x}) + \sigma(\check{c}, \check{x})\check{h}] \cdot V_x(\check{x}) \\ &\quad + \frac{1}{2} \text{trace} [\sigma(\check{c}, \check{x})' V_{xx}(\check{x}) \sigma(\check{c}, \check{x})], \end{aligned} \tag{32}$$

which involves only the  $\check{x}$  component of the state vector and not  $\check{z}$ .<sup>17</sup>

A Bellman-Isaacs condition renders inconsequential the order of action taken in the recursive game. The Bellman-Isaacs condition requires:

**Assumption 7.1.** *The value function  $V$  satisfies*

$$\begin{aligned} \delta V(\check{x}) &= \max_{\check{c} \in \check{C}} \min_{\check{h}} U(\check{c}, \check{x}) + \frac{\theta}{2} \check{h} \cdot \check{h} + [\mu(\check{c}, \check{x}) + \sigma(\check{c}, \check{x})\check{h}] \cdot V_x(\check{x}) \\ &\quad + \frac{1}{2} \text{trace} [\sigma(\check{c}, \check{x})' V_{xx}(\check{x}) \sigma(\check{c}, \check{x})] \\ &= \min_{\check{h}} \max_{\check{c} \in \check{C}} U(\check{c}, \check{x}) + \frac{\theta}{2} \check{h} \cdot \check{h} + [\mu(\check{c}, \check{x}) + \sigma(\check{c}, \check{x})\check{h}] \cdot V_x(\check{x}) \\ &\quad + \frac{1}{2} \text{trace} [\sigma(\check{c}, \check{x})' V_{xx}(\check{x}) \sigma(\check{c}, \check{x})] \end{aligned}$$

Appendix D describes three ways to verify this Bellman-Isaacs condition. The infinite-horizon counterpart to the result of Fleming and Souganidis (1989) asserts that the Bellman-Isaacs condition implies assumption 5.5 and hence  $\check{V}(\theta) = V(x_0)$  because  $z$  is initialized at unity.

---

as would be required for the HJB equation in (32) below to possess classical solutions. However Fleming and Souganidis (1989) prove that the value function satisfies the HJB equation in a weaker *viscosity* sense. Viscosity solutions are often needed when it is feasible and sometimes desirable to set the control  $\check{c}$  so that  $\sigma(\check{c}, \check{x})$  has lower rank than  $d$ , which is the dimension of the Brownian motion.

<sup>17</sup>We can construct another differential game for which  $V$  is the value function replacing  $dB_t$  by  $h_t dt + dB_t$  in the evolution equation instead of introducing a martingale. In this way we would perturb the process rather than the probability distribution. While this approach can be motivated using Girsanov's Theorem, some subtle differences between the resulting perturbation game and the martingale game arise because the history of  $\check{B}_t = \int_0^t h_u du + B_t$  can generate either a smaller or a larger filtration than that of the Brownian motion  $B$ . When it generates a smaller sigma algebra, we would be compelled to solve a combined control and filtering problem if we think of  $\check{B}$  as the generating the information available to the decision maker. If  $\check{B}$  generates a larger information set, then we are compelled to consider weak solutions to the stochastic differential equations that underlie the decision problem. Instead of extensively developing this alternative interpretation of  $V$  (as we did in an earlier draft), we simply think of the partial differential equation (32) as a means of simplifying the solution to the martingale problem.

## 7.1 A representation of $z^*$

One way to represent the worst-case martingale  $z^*$  in the recursive penalty game opens a natural transition to the risk-sensitive ordinary control problem whose HJB equation is (13). The minimizing player's decision rule is  $\check{h} = \alpha_h(\check{x})$ , where

$$\alpha_h(\check{x}) = -\frac{1}{\theta} \sigma^*(\check{x})' V_x(\check{x}) \quad (33)$$

and  $\sigma^*(\check{x}) \equiv \sigma^*(\alpha_c(\check{x}), \check{x})$ . Suppose that  $V(\check{x})$  is twice continuously differentiable. Applying the formula on page 226 of Revuz and Yor (1994), form the martingale:

$$z_t^* = \exp \left( -\frac{1}{\theta} [V(x_t) - V(x_0)] - \int_0^t w(x_u) du \right),$$

where  $w$  is constructed to ensure that  $z^*$  has a zero drift. The worst case distribution assigns more weight to bad states as measured by an exponential adjustment to the value function. This representation leads directly to the risk-sensitive control problem that we take up in the next subsection.

## 7.2 Risk sensitivity revisited

The HJB equation for the recursive, risk-sensitive control problem is obtained by substituting the solution (33) for  $h$  into the partial differential equation (32):

$$\begin{aligned} \delta V(\check{x}) &= \max_{\check{c} \in \check{C}} \min_{\check{h}} U(\check{c}, \check{x}) + \frac{\theta}{2} \check{h} \cdot \check{h} + [\mu(\check{c}, \check{x}) + \sigma(\check{c}, \check{x}) \check{h}] \cdot V_x(\check{x}) \\ &\quad + \frac{1}{2} \text{trace} [\sigma(\check{c}, \check{x})' V_{xx}(\check{x}) \sigma(\check{c}, \check{x})] \\ &= \max_{\check{c} \in \check{C}} U(\check{c}, \check{x}) + \mu(\check{c}, \check{x}) \cdot V_x(\check{x}) \\ &\quad + \frac{1}{2} \text{trace} [\sigma(\check{c}, \check{x})' V_{xx}(\check{x}) \sigma(\check{c}, \check{x})] \\ &\quad - \frac{1}{2\theta} V_x(\check{x})' \sigma(\check{c}, \check{x}) \sigma(\check{c}, \check{x})' V_x(\check{x}) \end{aligned} \quad (34)$$

The value function  $V$  for the robust penalty problem is also the value function for the risk sensitive control problem of section 3.2. The *risk sensitive* interpretation excludes worries about misspecified dynamics and instead enhances the control objective with aversion to risk in a way captured by the local variance of the continuation value. While mathematically related to the situation discussed in James (1992) (see pages 403 and 404), the presence of discounting in our setup compels us to use a recursive representation of the objective of the decision-maker.

In light of this connection between robust control and risk-sensitive control, it is not surprising that the penalty preference ordering that we developed in section 5.3 is equivalent

to a risk-sensitive version of the stochastic differential utility studied by Duffie and Epstein (1992). Using results from Schroder and Skiadas (1999), Skiadas (2001) has shown this formally.

The equivalence of the robustness-penalty preference order with one coming from a risk-adjustment of the continuation value obviously provides no guidance about which interpretation we should prefer. That a given preference order can be motivated in two ways does not inform us about which of them is more attractive. But in an application to asset pricing, Anderson, Hansen, and Sargent (2003) have shown how the robustness motivation would lead a calibrator to think differently about the parameter  $\theta$  than the risk motivation.<sup>18</sup>

## 8 Sequential timing protocol for a constraint formulation

Section 7 showed how to make penalty problem 5.2 recursive by adopting a sequential timing protocol. Now we show how to make the constraint problem 5.3 recursive. Because the value of the date zero constraint problem depends on the magnitude of the entropy constraint, we add the continuation value of entropy as a state variable. Instead of a value function  $V$  that depends only on the state  $x$ , we use a value function  $K$  that also depends on continuation entropy, denoted  $r$ .

### 8.1 An HJB equation for a constraint game

Our strategy is to use the link between the value functions for the penalty and constraint problems asserted in claims 5.4 and 5.6, then to deduce from the HJB equation (31) a partial differential equation that can be interpreted as the HJB equation for another zero-sum two-player game with additional states and controls. By construction, the new game has a sequential timing protocol and will have the same equilibrium outcome and representation as game (31). Until now, we have suppressed the dependence of  $V$  on  $\theta$  in our notation for the value function  $V$ . Because this dependence is central, we now denote it explicitly.

### 8.2 Another value function

Claim 5.4 showed how to construct the date zero value function for the constraint problem from the penalty problem via Legendre transform. We use this same transform over time to construct a new value function  $K$ :

$$K(\tilde{x}, \tilde{r}) = \max_{\theta \geq 0} V(\tilde{x}, \theta) - \tilde{r}\theta \quad (35)$$

---

<sup>18</sup>The link between the preference orders would vanish if we limited the concerns about model misspecification to some components of the vector Brownian motion. In Wang (2001)'s axiomatic treatment, the preferences are defined over both the approximating model and the family of perturbed models. Both can vary. By limiting the family of perturbed models, we can break the link with recursive utility theory.

that is related to  $\tilde{K}$  by

$$\tilde{K}(\tilde{r}) = K(\tilde{x}, \tilde{r})$$

provided that  $\tilde{x}$  is equal to the date zero state  $x_0$ ,  $\tilde{r}$  is used for the initial entropy constraint, and  $\tilde{z} = 1$ . We also assume that the Bellman-Isaacs condition is satisfied, so that the inverse Legendre transform can be applied:

$$V(\tilde{x}, \theta) = \min_{\tilde{r} \geq 0} K(\tilde{x}, \tilde{r}) + \tilde{r}\theta. \quad (36)$$

When  $K$  and  $V$  are related by the Legendre transforms (35) and (36), their derivatives are closely related, if they exist. We presume the smoothness needed to compute derivatives.

The HJB equation (31) that we derived for  $V$  held for each value of  $\theta$ . We consider the consequences of varying the pair  $(\tilde{x}, \theta)$ , as in the construction of  $V$ , or we consider varying the pair  $(\tilde{x}, \tilde{r})$ , as in the construction of  $K$ . We have

$$K_r = -\theta \quad \text{or} \quad V_\theta = \hat{r}.$$

For a fixed  $\tilde{x}$ , we can vary  $\tilde{r}$  by changing  $\theta$ , or conversely we can vary  $\theta$  by changing  $\tilde{r}$ . To construct a partial differential equation for  $K$  from (31), we will compute derivatives with respect to  $\tilde{r}$  that respect the constraint linking  $\tilde{r}$  and  $\theta$ .

For the optimized value of  $\tilde{r}$ , we have

$$\delta V = \delta(K + \theta\tilde{r}) = \delta K - \delta\tilde{r}K_r, \quad (37)$$

and

$$-\theta \left( \frac{\check{h} \cdot \check{h}}{2} \right) = K_r \left( \frac{\check{h} \cdot \check{h}}{2} \right). \quad (38)$$

By the implicit function theorem, holding  $\theta$  fixed:

$$\frac{\partial \tilde{r}}{\partial x} = -\frac{K_{xr}}{K_{rr}}.$$

Next we compute the derivatives of  $V$  that enter the partial differential equation (31) for  $V$ :

$$\begin{aligned} V_x &= K_x \\ V_{xx} &= K_{xx} + K_{rx} \frac{\partial \tilde{r}}{\partial x} \\ &= K_{xx} - \frac{K_{rx}K_{xr}}{K_{rr}}. \end{aligned} \quad (39)$$

Notice that

$$\begin{aligned} &\frac{1}{2} \text{trace} [\sigma(\check{c}, \check{x})' V_{xx}(\check{x}) \sigma(\check{c}, \check{x})] = \\ \min_{\check{g}} &\frac{1}{2} \text{trace} \left( \begin{bmatrix} \sigma(\check{c}, \check{x})' & \check{g} \end{bmatrix} \begin{bmatrix} K_{xx}(\check{x}, \tilde{r}) & K_{xr}(\check{x}, \tilde{r}) \\ K_{rx}(\check{x}, \tilde{r}) & K_{rr}(\check{x}, \tilde{r}) \end{bmatrix} \begin{bmatrix} \sigma(\check{c}, \check{x}) \\ \check{g}' \end{bmatrix} \right) \end{aligned} \quad (40)$$

where  $\check{g}$  is a column vector with the same dimension  $d$  as the Brownian motion. Substituting equations (37), (38), (39), and (40) into the partial differential equation (32) gives:

$$\begin{aligned} \delta K(\check{x}, \check{r}) &= \max_{\check{c} \in \check{C}} \min_{\check{h}, \check{g}} U(\check{c}, \check{x}) + [\mu(\check{c}, \check{x}) + \sigma(\check{c}, \check{x})\check{h}] \cdot K_x(\check{x}, \check{r}) + \left( \delta \check{r} - \frac{\check{h} \cdot \check{h}}{2} \right) \cdot K_r(\check{x}, \check{r}) \\ &\quad + \frac{1}{2} \text{trace} \left( \begin{bmatrix} \sigma(\check{c}, \check{x})' & \check{g} \end{bmatrix} \begin{bmatrix} K_{xx}(\check{x}, \check{r}) & K_{xr}(\check{x}, \check{r}) \\ K_{rx}(\check{x}, \check{r}) & K_{rr}(\check{x}, \check{r}) \end{bmatrix} \begin{bmatrix} \sigma(\check{c}, \check{x}) \\ \check{g}' \end{bmatrix} \right). \end{aligned} \quad (41)$$

The remainder of this section interprets  $\check{z}K(\check{x}, \check{r})$  as a value function for a recursive game in which  $\theta = \theta^* > \underline{\theta}$  is fixed over time. We have already seen how to characterize the state evolution for the recursive penalty differential game associated with a fixed  $\theta$ . The first-order condition for the maximization problem on the right side of (35) is

$$\check{r} = V_\theta(\check{x}, \theta^*). \quad (42)$$

We view this first-order condition as determining  $\check{r}$  for a *given*  $\theta^*$  and  $\check{x}$ . Then formula (42) implies that the evolution of  $r$  is fully determined by the equilibrium evolution of  $x$ . We refer to  $r$  as *continuation entropy*.

We denote the state evolution for the  $\theta^*$  differential game as:

$$dx_t = \mu^*(x_t, \theta^*)dt + \sigma^*(x_t, \theta^*)dB_t$$

### 8.3 Continuation entropy

We want to show that  $r$  evolves like continuation entropy. Recall formula (27) for the relative entropy of a nonnegative martingale:

$$\mathcal{R}^*(z) \doteq E \int_0^\infty \exp(-\delta t) z_t \frac{|h_t|^2}{2} dt.$$

Define a date  $t$  conditional counterpart as follows:

$$\mathcal{R}_t^*(z) = E \left[ \int_0^\infty \exp(-\delta u) \left( \frac{z_{t+u}}{z_t} \right) \frac{|h_{t+u}|^2}{2} du \middle| \mathcal{F}_t \right],$$

provided that  $z_t > 0$  and define  $\mathcal{R}_t^*(z)$  to be zero otherwise. This family of random variables induces the following recursion for  $\epsilon > 0$ :

$$z_t \mathcal{R}_t^*(z) = \exp(-\delta \epsilon) E \left[ z_{t+\epsilon} \mathcal{R}_{t+\epsilon}^*(z) \middle| \mathcal{F}_t \right] + E \left[ \int_0^\epsilon \exp(-\delta u) z_{t+u} \frac{|h_{t+u}|^2}{2} du \middle| \mathcal{F}_t \right].$$

Since  $z_t \mathcal{R}_t^*(z)$  is in the form of a risk neutral value of an asset with future *dividend*  $z_{t+u} \frac{|h_{t+u}|^2}{2}$ , its local mean or drift has the familiar formula:

$$\delta z_t \mathcal{R}_t^*(z) - z_t \frac{|h_t|^2}{2}.$$

To defend an interpretation of  $r_t$  as continuation entropy, we need to verify that this drift restriction is satisfied for  $r_t = \mathcal{R}_t^*(z)$ . Write the evolution for  $r_t$  as:

$$dr_t = \mu_r(x_t)dt + \sigma_r(x_t) \cdot dB_t,$$

and recall that

$$dz_t = z_t h_t \cdot dB_t.$$

Using Ito's formula for the drift of  $z_t r_t$ , the restriction that we want to verify is:

$$\check{z}\mu_r(\check{x}) + \check{z}\sigma_r(\check{x}) \cdot \check{h} = \delta\check{z}\check{r} - \check{z}\frac{|\check{h}|^2}{2}. \quad (43)$$

Given formula (42) and Ito's differential formula for a smooth function of a diffusion process, we have

$$\mu_r(\check{x}) = V_{\theta x}(\check{x}, \theta^*) \cdot \mu^*(\check{x}, \theta^*) + \frac{1}{2} \text{trace} [\sigma(\check{c}, \check{x})' V_{\theta xx}(\check{x}) \sigma(\check{c}, \check{x})]$$

and

$$\sigma_r(\check{x}) = V_{\theta x}(\check{x}, \theta^*) \sigma^*(\check{x}, \theta^*).$$

Recall that the worst case  $h_t$  is given by

$$h_t = -\frac{1}{\theta^*} \sigma^*(x_t, \theta^*)' V_x(x_t, \theta^*)$$

and thus

$$\frac{|h_t|^2}{2} = \left( \frac{1}{2\theta^{*2}} \right) V_x(\check{x})' \sigma(\check{c}, \check{x}) \sigma(\check{c}, \check{x})' V_x(\check{x}).$$

Restriction (43) can be verified by substituting our formulas for  $r_t$ ,  $h_t$ ,  $\mu_r$  and  $\sigma_r$ . The resulting equation is equivalent to that obtained by differentiating the HJB equation (34) with respect to  $\theta$ , justifying our interpretation of  $r_t$  as a continuation entropy.

## 8.4 Minimizing continuation entropy

Having defended a specific construction of continuation entropy that supports a constant value of  $\theta$ , we now describe a differential game that makes entropy an endogenous state variable. To formulate that game, we consider the inverse Legendre transform (36) from which we construct  $V$  from  $K$  by minimizing  $\check{r}$ . In the recursive version of the constraint game, the state variable  $r_t$  is the continuation entropy that at  $t$  remains available to allocate across states at future dates. At date  $t$ , continuation entropy is allocated via the minimization suggested by the inverse Legendre transform. We restrict the minimizing player to allocate future  $r_t$  across states that can be realized with positive probability, conditional on date  $t$  information.

### 8.4.1 Two state example

Before presenting the continuous-time formulation, consider a two-period example. Suppose that two states can be realized at date  $t + 1$ , namely  $\omega_1$  and  $\omega_2$ . Each state has probability one-half under an approximating model. The minimizing agent distorts these probabilities by assigning probability  $p_t$  to state  $\omega_1$ . The contribution to entropy coming from the distortion of the probabilities is the discrete state analogue of  $\int \log \left( \frac{dq_t}{dq_t^0} \right) dq_t$ , namely,

$$I(p_t) = p_t \log p_t + (1 - p_t) \log(1 - p_t) + \log 2.$$

The minimizing player also chooses continuation entropies for each of the two states that can be realized next period. Continuation entropies are discounted and averaged according to the distorted probabilities, so that we have:

$$r_t = I(p_t) + \exp(-\delta) [p_t r_{t+1}(\omega_1) + (1 - p_t) r_{t+1}(\omega_2)]. \quad (44)$$

Let  $U_t$  denote the current period utility for an exogenously given process for  $c_t$ , and let  $V_{t+1}(\omega, \theta)$  denote the next period value given state  $\omega$ . This function is concave in  $\theta$ . Construct  $V_t$  via backward induction:

$$V_t(\theta) = \min_{0 \leq p_{t+1} \leq 1} U_t + \theta I_t(p_t) + \exp(-\delta) [p_t V_{t+1}(\omega_1, \theta) + (1 - p_t) V_{t+1}(\omega_2, \theta)] \quad (45)$$

Compute the Legendre transforms:

$$\begin{aligned} K_t(\check{r}) &= \max_{\theta \geq 0} V_t(\theta) - \theta \check{r} \\ K_{t+1}(\check{r}, \omega) &= \max_{\theta \geq 0} V_{t+1}(\theta, \omega) - \theta \check{r} \end{aligned}$$

for  $\omega = \omega_1, \omega_2$ . Given  $\theta^*$ , let  $r_t$  be the solution to the inverse Legendre transform:

$$V_t(\theta^*) = \min_{\check{r} \geq 0} K_t(\check{r}) + \theta^* \check{r}.$$

Similarly, let  $r_{t+1}(\omega)$  be the solution to

$$V_{t+1}(\omega, \theta^*) = \min_{\check{r} \geq 0} K_{t+1}(\omega, \check{r}) + \theta^* \check{r}.$$

Substitute the inverse Legendre transforms into the simplified HJB equation (45):

$$\begin{aligned} V_t(\theta^*) &= \min_{0 \leq p_t \leq 1} U_t + \theta^* I_t(p_t) \\ &\quad + \exp(-\delta) \left( p_t \left[ \min_{\check{r}_1 \geq 0} K_{t+1}(\omega_1, \check{r}_1) + \theta^* \check{r}_1 \right] + (1 - p_t) \left[ \min_{\check{r}_2 \geq 0} K_{t+1}(\omega_2, \check{r}_2) + \theta^* \check{r}_2 \right] \right) \\ &= \min_{0 \leq p_t \leq 1, \check{r}_1 \geq 0, \check{r}_2 \geq 0} U_t + \theta^* (I_t(p_t) + \exp(-\delta) [p_t \check{r}_1 + (1 - p_t) \check{r}_2]) \end{aligned}$$

$$+ \exp(-\delta) [p_t K_{t+1}(\omega_1, \check{r}_1) + (1 - p_t) K_{t+1}(\omega_2, \check{r}_2)].$$

Thus,

$$\begin{aligned} K_t(r_t) &= V_t(\theta^*) - \theta^* r_t \\ &= \min_{0 \leq p_t \leq 1, \check{r}_1 \geq 0, \check{r}_2 \geq 0} \max_{\theta \geq 0} U_t + \theta (I_t(p_t) + \exp(-\delta) [p_t \check{r}_1 + (1 - p_t) \check{r}_2] - r_t) \\ &\quad + \exp(-\delta) [p_t K_{t+1}(\omega_1, \check{r}_1) + (1 - p_t) K_{t+1}(\omega_2, \check{r}_2)]. \end{aligned}$$

Since the solution is  $\theta = \theta^* > 0$ , at this value of  $\theta$  the entropy constraint (44) must be satisfied and

$$K_t(r_t) = \min_{0 \leq p_t \leq 1, \check{r}_1 \geq 0, \check{r}_2 \geq 0} U_t + \exp(-\delta) [p_t K_{t+1}(\omega_1, \check{r}_1) + (1 - p_t) K_{t+1}(\omega_2, \check{r}_2)].$$

By construction, the solution for  $\check{r}_j$  is  $r_{t+1}(\omega_j)$  defined earlier. The recursive implementation presumes that the continuation entropies  $r_{t+1}(\omega_j)$  are chosen at date  $t$  prior to the realization of  $\omega$ .

When we allow the decision maker to choose the control  $c_t$ , this construction requires that we can freely change orders of maximization and minimization as in our previous analysis.

#### 8.4.2 Continuous-time formulation

In a continuous-time formulation, we allocate the stochastic differential of entropy subject to the constraint that the current entropy is  $r_t$ . The increment to  $r$  is determined via the stochastic differential equation:<sup>19</sup>

$$dr_t = \left( \delta r_t - \frac{|h_t|^2}{2} - g_t \cdot h_t \right) dt + g_t \cdot dB_t.$$

This evolution for  $r$  implies that

$$d(z_t r_t) = \left( \delta z_t r_t - z_t \frac{|h_t|^2}{2} \right) dt + z_t (r_t h_t + g_t) dB_t$$

which has the requisite drift to interpret  $r_t$  as continuation entropy.

The minimizing agent not only picks  $h_t$  but also chooses  $g_t$  to allocate entropy over the next instant. The process  $g$  thus becomes a control vector for allocating continuation entropy across the various future states. In formulating the continuous-time game, we thus add a state  $r_t$  and a control  $g_t$ . With these added states, the differential game has a value function  $\hat{z}K(\hat{x}, \hat{r})$ , where  $K$  satisfies the HJB equation (41).

We have deduced this new partial differential equation partly to help us understand senses in which the constrained problem is or is not time consistent. Since  $r_t$  evolves as an exact function of  $x_t$ , it is more efficient to compute  $V$  and to use this value function to infer the optimal control law and the implied state evolution. In the next section, however, we use the recursive constraint formulation to address some interesting issues raised by Epstein and Schneider (2004).

---

<sup>19</sup>The process is stopped if  $r_t$  hits the zero boundary. Once zero is hit, the continuation entropy remains at zero. In many circumstances, the zero boundary will never be hit.

## 9 A recursive multiple priors formulation

Taking continuation entropy as a state variable is a convenient way to restrict the models entertained at time  $t$  by the minimizing player in the recursive version of constraint game. Suppose instead that at date  $t$  the decision maker retains the date zero family of probability models without imposing additional restrictions or freezing a state variable like continuation entropy. That would allow the minimizing decision maker at date  $t$  to reassign probabilities of events that have already been realized and events that cannot possibly be realized given current information. The minimizing decision maker would take advantage of that opportunity to alter the worst-case probability distribution at date  $t$  in a way that makes the specification of prior probability distributions of section 5 induce dynamic inconsistency in a sense formalized by Epstein and Schneider (2004). They characterize families of prior distributions that satisfy a rectangularity criterion that shields the decision maker from what they call “dynamic inconsistency”. In this section, we discuss how Epstein and Schneider’s notion of dynamic inconsistency would apply to our setting, show that their proposal for attaining consistency by minimally enlarging an original set of priors to be rectangular will not work for us, then propose our own way of making priors rectangular in a way that leaves the rest of our analysis intact.

Consider the martingale formulation of the date zero entropy constraint:

$$E \int_0^\infty \exp(-\delta u) z_u \frac{|h_u|^2}{2} du \leq \eta \quad (46)$$

where

$$dz_t = z_t h_t \cdot dB_t.$$

The component of entropy that constrains our date  $t$  decision-maker is:

$$r_t = \frac{1}{z_t} E \left( \int_0^\infty z_{t+u} \frac{|h_{t+u}|^2}{2} du \middle| \mathcal{F}_t \right)$$

in states in which  $z_t > 0$ . We rewrite (46) as:

$$E \int_0^t \exp(-\delta u) z_u \frac{|h_u|^2}{2} du + \exp(-\delta t) E z_t r_t \leq \eta.$$

To illuminate the nature of dynamic inconsistency, we begin by noting that the time 0 constraint imposes essentially no restriction on  $r_t$ . Consider a date  $t$  event that has probability strictly less than one conditioned on date zero information. Let  $y$  be a random variable that is equal to zero on the event and equal to the reciprocal of the probability on the complement of the event. Thus,  $y$  is a nonnegative, bounded random variable with expectation equal to unity. Construct a  $z_u = E(y|\mathcal{F}_u)$ . Then  $z$  is a bounded nonnegative martingale with finite entropy and  $z_u = y$  for  $u \geq t$ . In particular  $z_t$  is zero on the date  $t$  event used to construct  $y$ . By shrinking the date  $t$  event to have arbitrarily small probability, we can bring the bound arbitrarily close to unity and entropy arbitrarily close to zero. Thus, for date

$t$  events with sufficiently small probability, the entropy constraint can be satisfied without restricting the magnitude of  $r_t$  on these events. This exercise isolates a justification for using continuation entropy as a state variable inherited at date  $t$ : fixing it eliminates any gains from readjusting distortions of probabilities assigned to uncertainties that were resolved in previous time periods

## 9.1 Epstein and Schneider’s proposal works poorly for us

If we insist on withdrawing an endogenous state variable like  $r_t$ , dynamic consistency can still be obtained by imposing restrictions on  $h_t$  for alternative dates and states. For instance, we could impose prior restrictions in the separable form

$$\frac{|h_t|^2}{2} \leq f_t$$

for each event realization and date  $t$ . Such a restriction is *rectangular* in the sense of Epstein and Schneider (2004). To preserve a subjective notion of prior distributions, Epstein and Schneider (2004) advocate making an original set of priors rectangular by enlarging it to the least extent possible. They suggest this approach in conjunction with entropy measures of the type used here, as well as other possible specifications. However, an  $f_t$  specified on any event that occurs with probability less than one is essentially unrestricted by the date zero entropy constraint. In continuous time, this follows because zero measure is assigned to any calendar date, but it also carries over to discrete time because continuation entropy remains unrestricted if we can adjust earlier distortions. Thus, for our application Epstein and Schneider’s way of achieving a rectangular specification through the mechanism fails to restrict prior distributions in an interesting way.<sup>20</sup>

## 9.2 A better way to impose rectangularity

There is an alternative way to make the priors rectangular that has trivial consequences for our analysis. The basic idea is to separate the choice of  $f_t$  from the choice of  $h_t$ , while imposing  $\frac{|h_t|^2}{2} \leq f_t$ . We then imagine that the process  $\{f_t : t \geq 0\}$  is chosen *ex ante* and adhered to. Conditioned on that commitment, the resulting problem has the recursive structure advocated by Epstein and Schneider (2004). The ability to exchange maximization and minimization is central to our construction.

From section 5, recall that

$$\tilde{K}(\tilde{r}) = \max_{\theta \geq 0} \tilde{V}(\theta) - \theta \tilde{r}.$$

We now rewrite the inner problem on the right side for a fixed  $\theta$ . Take the Bellman-Isaacs condition

$$zV(x) = \min_{h \in H} \max_{c \in C} E \int_0^\infty \exp(-\delta t) \left[ z_t U(c_t, x_t) + \theta z_t \frac{|h_t|^2}{2} \right] dt$$

---

<sup>20</sup>While Epstein and Schneider (2004) advocate rectangularization even for entropy-based constraints, they do not claim that it always gives rise to interesting restrictions.

with the evolution equations

$$\begin{aligned} dx_t &= \mu(c_t, x_t)dt + \sigma(c_t, x_t)dB_t \\ dz_t &= z_t h_t \cdot dB_t. \end{aligned} \tag{47}$$

Decompose the entropy constraint as:

$$\eta = E \int_0^\infty \exp(-\delta t) z_t f_t dt$$

where

$$f_t = \frac{|h_t|^2}{2}.$$

Rewrite the objective of the optimization problem as

$$\min_{f \in F} \min_{h \in H, \frac{|h_t|^2}{2} \leq f_t} \max_{c \in C} E \int_0^\infty \exp(-\delta t) [z_t U(c_t, x_t) + \theta z_t f_t] dt$$

subject to (47). In this formulation,  $F$  is the set of progressively measurable scalar processes that are nonnegative. We entertain the inequality

$$\frac{|h_t|^2}{2} \leq f_t$$

but in fact this constraint will always bind for the *a priori* optimized choice of  $f$ . The inner problem can now be written as:

$$\min_{h \in H, \frac{|h_t|^2}{2} \leq f_t} \max_{c \in C} E \int_0^\infty \exp(-\delta t) z_t U(c_t, x_t) dt$$

subject to (47). Provided that we can change orders of the min and max, this inner problem will have a rectangular specification of alternative models and be dynamically consistent in the sense of Epstein and Schneider (2004).

Although this construction avoids introducing continuation entropy as an endogenous state variable, it assumes a commitment to a process  $f$  that is computed *ex ante* by solving what is essentially a static optimization problem. That is,  $f$  is chosen by exploring its consequences for a dynamic implementation of the form envisioned by Epstein and Schneider (2004) and is not simply part of the exogenously *ex ante* given set of beliefs of the decision maker.<sup>21</sup> We can, however, imagine that at date zero, the decision maker accepts the sequence  $\{f_t : t \geq 0\}$  as part of a conditional preference formulation. This decision maker then has preferences of a type envisioned by Epstein and Schneider (2004).

While their concern about dynamic consistency leads Epstein and Schneider to express doubts about commitments to a constraint based on continuation entropy, they do not

---

<sup>21</sup>Notice that the Bayesian interpretation is also a trivial special case of a recursive multiple priors model.

examine what could lead a decision-maker to commit to a particular rectangular set of beliefs embodied in a specification of  $f$ .<sup>22</sup> If multiple priors truly are a statement of a decision maker's subjective beliefs, we think it is not appropriate to dismiss such beliefs on the grounds of dynamic inconsistency. Repairing that inconsistency through the enlargements necessary to induce rectangularity reduces the content of the original set of prior beliefs. In our context, this enlargement is immense, too immense to be interesting to us.

The reservations that we have expressed about the substantive importance of rectangularity notwithstanding, we agree that Epstein and Schneider's discussion of dynamic consistency opens up a useful discussion of the alternative possible forms of commitment that allow us to create dynamic models with multiple priors.<sup>23</sup>

## 10 Concluding remarks

Empirical studies in macroeconomics and finance typically assume a unique and explicitly specified dynamic statistical model. Concerns about model misspecification recognize that an unknown member of a set of alternative models might govern the data. But how should one specify those alternative models? With one parameter that measures the size of the set, robust control theory parsimoniously stipulates a set of alternative models with rich dynamics.<sup>24</sup> Robust control theory leaves those models only vaguely specified and obtains them by perturbing the decision maker's approximating model to let shocks feed back on state variables arbitrarily. Among other possibilities, this allows the approximating model to miss the serial correlation of exogenous variables and the dynamics of how those exogenous variables impinge on endogenous state variables.

We have delineated some formal connections that exist between various formulations of robust control theory and the max-min expected utility theory of Gilboa and Schmeidler (1989). Their theory deduces a set of models from a decision maker's underlying preferences over risky outcomes. In their theory, none of the decision maker's models has the special status that the approximating model has in robust control theory. To put Gilboa and Schmeidler's theory to work, an applied economist would have to impute a set of models to the decision makers in his model (unlike the situation in rational expectations models, where the decision maker's model would be an equilibrium outcome). A practical attraction of robust control theory is the way it allows an economist to take a single approximating model and from it manufacture a set of models that express a decision maker's ambiguity.

---

<sup>22</sup>Furthermore, an analogous skeptical observation about commitment pertains to Bayesian decision theory, where the decision maker commits to a specific prior distribution.

<sup>23</sup>In the second to last paragraph of their page 16, Epstein and Schneider (2004) seem also to express reservations about their enlargement procedure.

<sup>24</sup>Other formulations of robust control put more structure on the class of alternative models and this can have important consequences for decisions. See Onatski and Williams (2003) for one more structured formulation and Hansen and Sargent (2005b) for another. By including a hidden state vector and appropriately decomposing the density of next period's observables conditional on a history of signals, Hansen and Sargent (2005b) extend the approach of this paper to allow a decision maker to have multiple models and to seek robustness to the specification of a prior over them.

Hansen and Sargent (2003) exploit this feature of robust control to construct a multiple agent model in which a common approximating model plays the role that an equilibrium common model does in a rational expectations model.

We have used a particular notion of discounted entropy as a statistical measure of the discrepancy between models. It directs our decision maker's attention to models that are absolutely continuous with respect to his approximating model over finite intervals, but not absolutely continuous with respect to it over an infinite interval. This specification keeps the decision maker concerned about models that can be difficult to distinguish from the approximating model from a continuous record of observations on the state vector of a finite length. Via statistical detection error probabilities, Anderson, Hansen, and Sargent (2003) show how the penalty parameter or the constraint parameter in the robust control problems can be used to identify a set of perturbed models that are difficult to distinguish statistically from the approximating model in light of a continuous record of finite length  $T$  of observations on  $x_t$ .

Finally, we have made extensive use of martingales to represent perturbed models. Hansen and Sargent (2005a) and Hansen and Sargent (2005b) use such martingales to pose robust control and estimation problems in Markov decision problems where some of the state variables are hidden.

## A Cast of characters

This appendix sets out the following list of objects and conventions that make repeated appearances in our analysis.

1. Probability spaces
  - (a) A probability space associated with a Brownian motion  $B$  that is used to define an approximating model and a set of alternative models.
  - (b) A probability space over continuous functions of time induced by history of the Brownian motion  $B$  in part 1a and used to define an approximating model.
  - (c) A set of alternative probability distributions induced by  $B$  and used to define a set alternative models.
2. Ordinary (single-agent) control problems
  - (a) A benchmark optimal control problem defined on space 1a.
  - (b) A benchmark decision problem defined on the probability space induced by  $B$ .
  - (c) A risk-sensitive problem defined on space 1a.
  - (d) Alternative Bayesian (benchmark problems) defined on the spaces in 1c.
3. Representations of alternative models
  - (a) As nonnegative martingales with unit expectation the probability space 1a.
  - (b) As alternative induced distributions as in 1c.
4. Restrictions on sets of alternative models
  - (a) An implicit restriction embedded in a nonnegative penalty parameter  $\theta$ .
  - (b) A constraint on relative entropy, a measure of model discrepancy.
5. Representations of relative entropy
  - (a) Time 0 (nonsequential): discounted expected log likelihood ratio of an approximating model  $q^0$  to an alternative model  $q$  drawn from the set 1c.
  - (b) Time 0 (nonsequential): a function of a martingale defined on the probability space 1a.
  - (c) Recursive: as a solution of either of a differential equations defined in terms of  $B$ .
6. Timing protocols for zero-sum two-player games
  - (a) Exchange of order of choice for maximizing and minimizing players.
  - (b) Under two-sided commitment at  $t = 0$ , both players choose processes for all time  $t \geq 0$ .
  - (c) With lack of commitment on two sides, both players choose sequentially.

## B Discounted entropy

Let  $Q$  be the set of all distributions that are absolutely continuous with respect to  $q^0$  over finite intervals. This set is convex. For  $q \in Q$ , let

$$\tilde{\mathcal{R}}(q) \doteq \delta \int_0^\infty \exp(-\delta t) \left[ \int \log \left( \frac{dq_t}{dq_t^0} \right) dq_t \right] dt,$$

which may be infinite for some  $q \in Q$ .

**Claim B.1.**  $\tilde{\mathcal{R}}$  is convex on  $Q$ .

*Proof.* Since  $q \in Q$  is absolutely continuous with respect to  $q^0$  over finite intervals, we can construct likelihood ratios for finite histories at any calendar date  $t$ . Form  $\tilde{\Omega} = \Omega^* \times \mathbb{R}^+$  where  $\mathbb{R}^+$  is the nonnegative real line. Form the corresponding sigma algebra  $\tilde{\mathcal{F}}$  as the smallest sigma algebra containing  $\mathcal{F}_t^* \otimes \mathcal{B}_t$  for any  $t$  where  $\mathcal{B}_t$  is the collection of Borel sets in  $[0, t]$ ; and form  $\tilde{q}$  as the product measure  $q$  with an exponential distribution with density  $\delta \exp(-\delta t)$  for any  $q \in Q$ . Notice that  $\tilde{q}$  is a probability distribution and  $\tilde{\mathcal{R}}(q)$  is the relative entropy of  $\tilde{q}$  with respect to  $\tilde{q}^0$ :

$$\tilde{\mathcal{R}}(q) = \int \log \left( \frac{d\tilde{q}}{d\tilde{q}^0} \right) d\tilde{q}.$$

Form two measures  $\tilde{q}^1$  and  $\tilde{q}^2$  as the product of  $q^1$  and  $q^2$  with an exponential distribution with parameter  $\delta$ . Then a convex combination of  $\tilde{q}^1$  and  $\tilde{q}^2$  is given by the product of the corresponding convex combination of  $q^1$  and  $q^2$  with the same exponential distribution. Relative entropy is well known to be convex in the probability measure  $\tilde{q}$  (e.g. see Dupuis and Ellis (1997)), and hence  $\tilde{\mathcal{R}}$  is convex in  $q$ .  $\square$

Recall that associated with any probability measure  $q$  that is absolutely continuous with respect to  $q^0$  over finite intervals is a nonnegative martingale  $z$  defined on  $(\Omega, \mathcal{F}, P)$  with a unit expectation. This martingale satisfies the integral equation:

$$z_t = 1 + \int_0^t z_u h_u dB_u. \tag{48}$$

**Claim B.2.** Suppose that  $q_t$  is absolutely continuous with respect to  $q_t^0$  for all  $0 < t < \infty$ . Let  $z$  be the corresponding nonnegative martingale on  $(\Omega, \mathcal{F}, P)$ . Then

$$E z_t \mathbf{1}_{\{\int_0^t |h_s|^2 ds < \infty\}} = 1.$$

Moreover,

$$\int \log \frac{dq_t}{dq_t^0} dq_t = \frac{1}{2} E \int_0^t z_s |h_s|^2 ds.$$

*Proof.* Consider first the claim that

$$E z_t \mathbf{1}_{\left\{\int_0^t |h_s|^2 ds < \infty\right\}} = 1,$$

The martingale  $z$  satisfies the stochastic differential equation:

$$dz_t = z_t h_t dB_t$$

with initial condition  $z_0 = 1$ . Construct an increasing sequence of stopping times  $\{\tau_n : n \geq 1\}$  where  $\tau_n \doteq \inf\{t : z_t = \frac{1}{n}\}$  and let  $\tau = \lim_n \tau_n$ . The limiting stopping time can be infinite. Then  $z_t = 0$  for  $t \geq \tau$  and

$$z_t = z_{t \wedge \tau}$$

Form:

$$z_t^n = z_{t \wedge \tau_n}$$

which is nonnegative martingale satisfying:

$$dz_t^n = z_t^n h_t^n dB_t$$

where  $h_t^n = h_t$  if  $0 < t < \tau_n$  and  $h_t^n = 0$  if  $t \geq \tau_n$ . Then

$$P \left\{ \int_0^t |h_s^n|^2 (z_s^n)^2 ds < \infty \right\} = 1$$

and hence

$$P \left\{ \int_0^t |h_s^n|^2 ds < \infty \right\} = P \left\{ \int_0^{t \wedge \tau_n} |h_s|^2 ds < \infty \right\} = 1.$$

Taking limits as  $n$  gets large,

$$P \left\{ \int_0^{t \wedge \tau} |h_s|^2 ds < \infty \right\} = 1.$$

While it is possible that  $\tau < \infty$  with positive  $P$  probability, as argued by Kabanov, Lipcer, and Sirjaev (1979)

$$\int z_t \mathbf{1}_{\{\tau < \infty\}} dP = \int_{\{z_t=0, t < \infty\}} z_t dP = 0.$$

Therefore,

$$E z_t \mathbf{1}_{\left\{\int_0^t |h_s|^2 ds < \infty\right\}} = E z_t \mathbf{1}_{\left\{\int_0^{t \wedge \tau} |h_s|^2 ds < \infty, \tau = \infty\right\}} + E z_t \mathbf{1}_{\left\{\int_0^t |h_s|^2 ds < \infty, \tau < \infty\right\}} = 1.$$

Consider next the claim that

$$\int \log \frac{dq_t}{dq_t^0} dq_t = E \int_0^t z_s |h_s|^2 ds.$$

We first suppose that

$$E \int_0^t z_s |h_s|^2 ds < \infty. \quad (49)$$

We will subsequently show that this condition is satisfied when  $\tilde{\mathcal{R}}(q) < \infty$ . Use the martingale  $z$  to construct a new probability measure  $\tilde{P}$  on  $(\Omega, \mathcal{F})$ . Then from the Girsanov Theorem [see Theorem 6.2 of Liptser and Shiryaev (2000)]

$$\tilde{B}_t = B_t - \int_0^t h_s ds$$

is a Brownian motion with respect to the filtration  $\{\mathcal{F}_t : t \geq 0\}$ . Moreover,

$$\tilde{E} \int_0^t |h_s|^2 ds = E \int_0^t z_s |h_s|^2 ds.$$

Write

$$\log z_t = \int_0^t h_s \cdot dB_s - \frac{1}{2} \int_0^t |h_s|^2 ds = \int_0^t h_s \cdot d\tilde{B}_s + \frac{1}{2} \int_0^t |h_s|^2 ds.$$

which is well defined under the  $\tilde{P}$  probability. Moreover,

$$\tilde{E} \int_0^t h_s \cdot d\tilde{B}_s = 0$$

and hence

$$\tilde{E} \log z_t = \frac{1}{2} \tilde{E} \int_0^t |h_s|^2 ds = \frac{1}{2} E \int_0^t z_s |h_s|^2 ds,$$

which is the desired equality. In particular, we have proved that  $\int \log \frac{dq_t}{dq_t^0} dq_t$  is finite.

Next we suppose that

$$\int \log \frac{dq_t}{dq_t^0} dq_t < \infty,$$

which will hold when  $\tilde{\mathcal{R}}(q) < \infty$ . Then Lemma 2.6 from Föllmer (1985) insures that

$$\frac{1}{2} \tilde{E} \int_0^t |h_s|^2 ds \leq \int \log \frac{dq_t}{dq_t^0} dq_t.$$

Föllmer's result is directly applicable because  $\int \log \frac{dq_t}{dq_t^0} dq_t$  is the same as the relative entropy of  $\tilde{P}_t$  with respect to  $P_t$  where  $\tilde{P}_t$  is the restriction of  $\tilde{P}$  to events in  $\mathcal{F}_t$  and  $P_t$  is defined similarly. As a consequence, (49) is satisfied and the desired equality follows from our previous argument.

Finally, notice that  $\frac{1}{2} \tilde{E} \int_0^t |h_s|^2 ds$  is infinite if, and only if  $\int \log \frac{dq_t}{dq_t^0} dq_t$  is infinite.  $\square$

**Claim B.3.** For  $q \in Q$ , let  $z$  be the nonnegative martingale associated with  $q$  and let  $h$  be the progressively measurable process satisfying (48). Then

$$\tilde{\mathcal{R}}(q) = \frac{1}{2} E \left[ \int_0^\infty \exp(-\delta t) z_t |h_t|^2 dt \right]$$

*Proof.* The conclusion follows from:

$$\begin{aligned} \tilde{\mathcal{R}}(q) &= \delta \int_0^\infty \exp(-\delta t) \int \log \left( \frac{dq_t}{dq_t^0} \right) dq_t dt \\ &= \frac{\delta}{2} E \left[ \int_0^\infty \exp(-\delta t) \int_0^t z_u |h_u|^2 du dt \right] \\ &= \frac{1}{2} E \left[ \int_0^\infty \exp(-\delta t) z_t |h_t|^2 dt \right] \end{aligned}$$

where the second equality follows from B.2 and the third from integrating by parts.  $\square$

This justifies our definition of entropy for nonnegative martingales:

$$\mathcal{R}(z) = \frac{1}{2} E \left[ \int_0^\infty \exp(-\delta t) z_t |h_t|^2 dt \right].$$

## C Absolute continuity of solutions

In this appendix we show how to verify that the solution for  $z$  from the martingale robust control problem is in a fact a martingale and not just a local martingale. Our approach to studying absolute continuity and verifying that the Markov perfect equilibrium  $z$  is a martingale differs from the perhaps more familiar use of a Novikov or Kazamaki condition.<sup>25</sup>

Consider two distinct stochastic differential equations. One is the Markov solution to the penalty robust control problem.

$$\begin{aligned} dx_t^* &= \mu^*(x_t^*) dt + \sigma^*(x_t^*) dB_t \\ dz_t^* &= z_t^* \alpha_h(x_t^*) dB_t. \end{aligned} \tag{50}$$

where  $\mu^*(\tilde{x}) = \mu(\alpha_c(\tilde{x}), \tilde{x})$ ,  $\sigma^*(\tilde{x}) = \sigma(\alpha_c(\tilde{x}), \tilde{x})$  and where  $\alpha_c$  and  $\alpha_h$  are the solutions from the penalty robust control problem. Notice that the equation for the evolution of  $x_t^*$  is autonomous (it does not depend on  $z_t^*$ ). Let a strong solution to this equation system be:

$$x_t^* = \Phi_t^*(B).$$

Consider a second stochastic differential equation:

$$d\hat{x}_t = \mu^*(\hat{x}_t) dt + \sigma^*(\hat{x}_t) \left[ \alpha_h(\hat{x}_t) dt + d\hat{B}_t \right] \tag{51}$$

---

<sup>25</sup>We construct two well defined Markov processes and verify absolute continuity. Application of the Novikov or Kazamaki conditions entails imposing extra moment conditions on the objects used to construct the local martingale  $z$ .

In verifying that this state equation has a solution, we are free to examine weak solutions provided that  $\hat{\mathcal{F}}_t$  is generated by current and past  $\hat{x}_t$  and  $\hat{B}$  does not generate a larger filtration than  $\hat{x}$ .

The equilibrium outcomes  $x^*$  and  $\hat{x}$  for the two stochastic differential equations thus induce two distributions for  $x$ . We next study how these distributions are related. We will discuss how to check for absolute continuity along finite intervals for induced distributions associated with these models. When the models satisfy absolute continuity over finite intervals, it will automatically follow that the equilibrium process  $z^*$  is a martingale.

## C.1 Comparing models of $B$

We propose the following method to transform a strong solution to (50) into a possibly weak solution to (51). Begin with a Brownian motion  $\hat{B}$  defined on a probability space with probability measure  $\hat{P}$ . Consider the recursive solution:

$$\begin{aligned}\hat{x}_t &= \Phi_t^*(B) \\ B_t &= \hat{B}_t + \int_0^t \alpha_h(\hat{x}_u) du.\end{aligned}$$

We look for solutions in which  $\mathcal{F}_t$  is generated by current and past values of  $B$  (not  $\hat{B}$ ). We call this a recursion because  $B$  is itself constructed from past values of  $B$  and  $\hat{B}$ . The stochastic differential equation associated with this recursion is (51).

To establish the absolute continuity of the distribution induced by  $B$  with respect to Wiener measure  $q^0$  it suffices to verify that for each  $t$

$$\hat{E} \int_0^t |\alpha_h(\hat{x}_u)|^2 du < \infty$$

and hence

$$\hat{P} \left\{ \int_0^t |\alpha_h(\hat{x}_u)|^2 du < \infty \right\} = 1. \quad (52)$$

It follows from Theorem 7.5 of Liptser and Shiryaev (2000) that the probability distribution induced by  $B$  under the solution to the perturbed problem is absolutely continuous with respect to Wiener measure  $q^0$ . To explore directly the weaker relation (52) further, recall that

$$\alpha_h(\tilde{x}) = -\frac{1}{\theta} \sigma^*(\tilde{x})' V_x(\tilde{x}).$$

Provided that  $\sigma^*$  and  $V_x$  are continuous in  $\tilde{x}$  and that  $x$  does not explode in finite time, this relation follows immediately.

## C.2 Comparing generators

Another strategy for checking absolute continuity is to follow the approach of Kunita (1969), who provides characterizations of absolute continuity and equivalence of Markov models

through restrictions on the generators of the processes. Since the models for  $x^*$  and  $\hat{x}$  are Markov diffusion processes, we can apply these characterizations provided that we include  $B$  as part of the state vector. Abstracting from boundary behavior, Kunita (1969) requires a common diffusion matrix, which can be singular. The differences in the drift vector are restricted to be in the range of the common diffusion matrix. These restrictions are satisfied in our application.

### C.3 Verifying $z^*$ is a martingale

We apply our demonstration of absolute continuity to reconsider the super martingale  $z^*$ . Let  $\kappa_t$  denote the Radon-Nikodym derivative for the two models of  $B$ . Conjecture that

$$z_t^* = \kappa_t(B).$$

By construction,  $z^*$  is a nonnegative martingale defined on  $(\Omega, \mathcal{F}, P)$ . Moreover, it is the unique solution to the stochastic differential equation (50) subject to the initial condition  $z_0^* = 1$ . See Theorem 7.6 of Liptser and Shiryaev (2000).

## D Three ways to verify the Bellman-Isaacs condition

This appendix describes three alternative conditions that are sufficient to verify the Bellman-Isaacs condition embraced in Assumption 7.1.<sup>26</sup> The ability to exchange orders of extremization in the recursive game implies that the orders of extremization can also be exchanged in the nonsequential game, as required in Assumption 5.5. As we shall now see, the exchange of order of extremization asserted in Assumption 7.1 can often be verified without knowing the value function  $S$ .

### D.1 No binding inequality restrictions

Suppose that there are no binding inequality restrictions on  $c$ . Then a justification for Assumption 7.1 can emerge from the first-order conditions for  $\check{c}$  and  $\check{h}$ . Define

$$\begin{aligned} \chi(\check{c}, \check{h}, \check{x}) \doteq & U(\check{c}, \check{x}) + \frac{\theta}{2} \check{h} \cdot \check{h} + [\mu(\check{c}, \check{x}) + \sigma(\check{c}, \check{x})\check{h}] \cdot S_x(\check{x}) \\ & + \frac{1}{2} \text{trace} [\sigma(\check{c}, \check{x})' S_{xx}(\check{x}) \sigma(\check{c}, \check{x})], \end{aligned} \quad (53)$$

---

<sup>26</sup>Fleming and Souganidis (1989) show that the freedom to exchange orders of maximization and minimization guarantees that equilibria of the nonsequential (i.e., choices under mutual commitment at date 0) and the recursive games (i.e., sequential choices by both agents) coincide.

and suppose that  $\chi$  is continuously differentiable in  $\check{c}$ . First, find a Markov perfect equilibrium by solving:

$$\begin{aligned}\frac{\partial \chi}{\partial c}(\check{c}^*, \check{h}^*, \check{x}) &= 0 \\ \frac{\partial \chi}{\partial h}(\check{c}^*, \check{h}^*, \check{x}) &= 0.\end{aligned}$$

In particular, the first-order conditions for  $\check{h}$  are:

$$\frac{\partial \chi}{\partial h}(\check{c}^*, \check{h}^*, \check{x}) = \theta \check{h}^* + \sigma(\check{c}^*, \check{x})' S_x(\check{x}) = 0.$$

If a unique solution exists and if it suffices for extremization, the Bellman-Isaacs condition is satisfied. This follows from the ‘‘chain rule.’’ Thus, suppose that the minimizing player goes first and computes  $\check{h}$  as a function of  $\check{x}$  and  $\check{c}$ :

$$\check{h}^* = -\frac{1}{\theta} \sigma(\check{c}, \check{x})' S_x(\check{x}) \tag{54}$$

Then the first-order conditions for the max player selecting  $\check{c}$  as a function of  $\check{x}$  are:

$$\frac{\partial \chi}{\partial c} + \frac{\partial h}{\partial c} \frac{\partial \chi}{\partial h} = 0$$

where  $\frac{\partial h}{\partial c}$  can be computed from the reaction function (54). Notice that the first-order conditions for the maximizing player are satisfied at the Markov perfect equilibrium. A similar argument can be made if the maximizing player chooses first.

## D.2 Separability

Consider next the case in which  $\sigma$  does not depend on the control. In this case the decision problems for  $\check{c}$  and  $\check{h}$  separate. For instance, from (54), we see that  $\check{h}$  does not react to  $\check{c}$  in the minimization of  $\check{h}$  conditioned on  $\check{c}$ . Even with binding constraints on  $\check{c}$ , the Bellman-Isaacs condition (Assumption 7.1) is satisfied, provided that a solution exists for  $\check{c}$ .

## D.3 Convexity

A third approach that uses results of Fan (1952) and Fan (1953) is based on the global shape properties of the objective. When we can reduce the choice set  $C$  to be a compact subset of a linear space, Fan (1952) can apply. Fan (1952) also requires that the set of conditional minimizers and maximizers be convex. We know from formula (54) that the minimizers of  $\chi(\check{c}, \cdot, \check{x})$  form a singleton set, which is convex for each  $\check{c}$  and  $\check{x}$ .<sup>27</sup> Suppose also that the

---

<sup>27</sup>Notice that provided  $\check{C}$  is compact, we can use (54) to specify a compact set that contains the entire family of minimizers for each  $\check{c}$  in  $\check{C}$  and a given  $\check{x}$ .

set of maximizers of  $\chi(\cdot, \check{h}, \check{x})$  is non-empty and convex for each  $\check{h}$  and  $\check{x}$ .<sup>28</sup> Then again the Bellman-Isaacs condition (Assumption 7.1) is satisfied. Finally Fan (1953) does not require that the set  $\check{C}$  be a subset of a linear space, but instead requires that  $\chi(\cdot, \check{h}, \check{x})$  be concave. By relaxing the linear space structure we can achieve compactness by adding points (say the point  $\infty$ ) to the control set, provided that we can extend  $\chi(\cdot, \check{h}, \check{x})$  to be upper semi-continuous. The extended control space must be a compact Hausdorff space. Provided that the additional points are not attained in optimization, we can apply Fan (1953) to verify Assumption 7.1.<sup>29</sup>

## E Recursive version of Stackelberg game and a Bayesian problem

### E.1 Recursive version of a Stackelberg game

We first change the timing protocol for decision-making, moving from the Markov perfect equilibrium that gives rise to a value function  $V$  to a date zero Stackelberg equilibrium with value function  $N$ . In the matrix manipulations that follow, state vectors and gradient vectors are treated as column vectors when they are pre-multiplied by matrices.

The value function  $V$  solves:

$$\begin{aligned} \delta V(\check{x}) &= \max_{\check{c} \in \check{C}} \min_{\check{h}} U(\check{c}, \check{x}) + \frac{\theta}{2} \check{h} \cdot \check{h} + [\mu(\check{c}, \check{x}) + \sigma(\check{c}, \check{x})\check{h}] \cdot V_x(\check{x}) \\ &\quad + \frac{1}{2} \text{trace} [\sigma(\check{c}, \check{x})' V_{xx}(\check{x}) \sigma(\check{c}, \check{x})] \end{aligned}$$

Associated with this value function are the first-order conditions for the controls:

$$\begin{aligned} \theta \check{h} + \sigma(\check{c}, \check{x})' \cdot V_x(\check{x}) &= 0 \\ \frac{\partial}{\partial \check{c}} \left( U(\check{c}, \check{x}) + [\mu(\check{c}, \check{x}) + \sigma(\check{c}, \check{x})\check{h}] \cdot V_x(\check{x}) + \frac{1}{2} \text{trace} [\sigma(\check{c}, \check{x})' V_{xx}(\check{x}) \sigma(\check{c}, \check{x})] \right) &= 0. \end{aligned}$$

Solving these first-order conditions gives the control laws  $h_t = \alpha(x_t)$  and  $c_t = \alpha_c(x_t)$ . Define  $\mu^*$  and  $\sigma^*$  such that the states evolve according to

$$dx_t = \mu^*(x_t)dt + \sigma^*(x_t)dB_t$$

after the two optimal controls are imposed. Associated with this recursive representation are processes  $h$  and  $c$  that can also be depicted as functions of the history of the underlying Brownian motion  $B$ .

<sup>28</sup>See Ekeland and Turnbull (1983) for a discussion of continuous time, deterministic control problems when the set of minimizers is not convex. They show that sometimes it is optimal to *chatter* between different controls as a way to imitate convexification in continuous time.

<sup>29</sup>Apply Theorem 2 of Fan (1953) to  $-\chi(\cdot, \cdot, \check{x})$ . This theorem does not require compactness of the choice set for  $\check{h}$ , only of the choice set for  $\check{c}$ . The theorem also does not require attainment when optimization is over the noncompact choice set. In our application, we can verify attainment directly.

When the Bellman-Isaacs condition is satisfied, Fleming and Souganidis (1989) provide a formal justification for an equivalent date zero Stackelberg solution in which the minimizing agent announces a decision process  $\{h_t : t \geq 0\}$  and the maximizing agent reacts by maximizing with respect to  $\{c_t : t \geq 0\}$ . We seek a recursive representation of this solution by using a big  $X$ , little  $x$  formulation. Posit a worst-case process for  $X_t$  of the form:

$$dX_t = \mu^*(X_t)dt + \sigma^*(X_t) [\alpha_h(X_t)dt + dB_t].$$

This big  $X$  process is designed so that it produces the same process for  $h_t = \alpha_h(X_t)$  that is implied by the Markov perfect equilibrium associated with the value function  $V$  when  $X_0 = x_0$ .

The big  $X$  process cannot be influenced by the maximizing agent, but little  $x$  can:

$$dx_t = \mu(c_t, x_t)dt + \sigma(c_t, x_t) [\alpha_h(X_t)dt + dB_t].$$

Combining the two state evolution equations, we have a Markov control problem faced by the maximizing agent. It gives rise to a value function  $N$  satisfying a HJB equation:

$$\begin{aligned} \delta N(\check{x}, \check{X}) &= \max_{\check{c} \in \check{C}} U(\check{c}, \check{x}) + \mu(\check{c}, \check{x}) \cdot N_x(\check{x}, \check{X}) + \mu^*(\check{x}) \cdot N_X(\check{X}, \check{X}) \\ &+ \frac{1}{2} \text{trace} \left( \begin{bmatrix} \sigma(\check{c}, \check{x})' & \sigma^*(\check{X})' \end{bmatrix} \begin{bmatrix} N_{xx}(\check{x}, \check{X}) & N_{xX}(\check{x}, \check{X}) \\ N_{Xx}(\check{x}, \check{X}) & N_{XX}(\check{x}, \check{X}) \end{bmatrix} \begin{bmatrix} \sigma(\check{c}, \check{x}) \\ \sigma^*(\check{X}) \end{bmatrix} \right) \\ &+ \alpha_h(\check{X}) \cdot \sigma(\check{c}, \check{x})' N_x(\check{x}, \check{X}) + \alpha_h(\check{X}) \cdot \sigma^*(\check{X})' N_X(\check{x}, \check{X}) \\ &+ \frac{\theta}{2} \alpha_h(\check{X}) \cdot \alpha_h(\check{X}). \end{aligned} \quad (55)$$

We want the outcome of this optimization problem to produce the same stochastic process for  $c$  ( $c_t$  as a function of current and past values of the Brownian motion  $B_t$ ) provided that  $X_0 = x_0$ . For this to happen, the value functions  $V$  and  $N$  must be closely related. Specifically,

$$\begin{aligned} N_x(\check{x}, \check{X})|_{\check{X}=\check{x}} &= V_x(\check{x}) \\ N_X(\check{x}, \check{X})|_{\check{X}=\check{x}} &= 0. \end{aligned} \quad (56)$$

The first restriction equates the co-state on little  $x$  with the implied co-state from the Markov perfect equilibrium along the equilibrium trajectory. The second restriction implies that the co-state vector for big  $X$  is zero along this same trajectory.

These restrictions on the first derivative, imply restrictions on the second derivative. Consider a perturbation of the form:

$$\check{x} + \mathbf{r}\nu, \quad \check{X} + \mathbf{r}\nu$$

for some scalar  $\mathbf{r}$  and some direction  $\nu$ . The directions that interest us are those in the range of  $\sigma^*(\check{X})$ , which are the directions that the Brownian motion can move the state to. Since (56) holds,

$$N_{xx}(\check{x}, \check{X})\nu + N_{xX}(\check{x}, \check{X})\nu|_{\check{X}=\check{x}} = V_{xx}(\check{x})\nu$$

$$N_{Xx}(\check{x}, \check{X})\nu + N_{XX}(\check{x}, \check{X})\nu|_{\check{X}=\check{x}} = 0.$$

From HJB (55), we could find a control law that expresses  $\check{c}$  as a function of  $\check{x}$  and  $\check{X}$ . We are only concerned, however, with  $\check{c}$  evaluated in the restricted domain  $\check{x} = \check{X}$ . Given the presumed restrictions on the first derivative and the derived restrictions on the second derivative, we can show that  $\check{c} = \alpha_c(\check{x})$  satisfies the first-order conditions for  $\check{c}$  provided on this restricted domain.

## E.2 Changing the objective

The value function for a Bayesian problem does not include a penalty term. In the recursive representation of the date zero Stackelberg problem, the penalty term is expressed completely in terms of big  $X$ . We now show how to adjust the value function  $L$  by solving a Lyapunov equation.

The function that we wish to compute solves:

$$L(\check{X}) = \frac{\theta}{2} E \int_0^\infty \exp(-\delta t) |\alpha_h(X_t)|^2 dt$$

subject to

$$dX_t = \mu^*(X_t)dt + \sigma^*(X_t) [\alpha_h(X_t)dt + dB_t].$$

where  $X_0 = \check{X}$ .

The value function  $L$  for this problem solves:

$$\begin{aligned} \delta L(\check{X}) &= \frac{\theta}{2} \alpha_h(\check{X}) \cdot \alpha_h(\check{X}) + \mu^*(\check{X}) \cdot L_X(\check{X}) \\ &+ \frac{1}{2} \text{trace} [\sigma^*(\check{X})' L_{XX}(\check{X}) \sigma^*(\check{X})] + \alpha_h(\check{X}) \cdot \sigma^*(\check{X})' L_x(\check{X}). \end{aligned} \quad (57)$$

## E.3 Bayesian value function

To construct a Bayesian value function we form:

$$W(\check{x}, \check{X}) = N(\check{x}, \check{X}) - L(\check{X}).$$

Given equations (55) and (57), the separable structure of  $W$  implies that it satisfies the HJB equation:

$$\begin{aligned} \delta W(\check{x}, \check{X}) &= \max_{\check{c} \in \check{C}} U(\check{c}, \check{x}) + \mu(\check{c}, \check{x}) \cdot W_x(\check{x}, \check{X}) + \mu^*(\check{x}) \cdot W_X(\check{X}, \check{X}) \\ &+ \frac{1}{2} \text{trace} \left( \begin{bmatrix} \sigma(\check{c}, \check{x})' & \sigma^*(\check{X})' \end{bmatrix} \begin{bmatrix} W_{xx}(\check{x}, \check{X}) & W_{xX}(\check{x}, \check{X}) \\ W_{Xx}(\check{x}, \check{X}) & W_{XX}(\check{x}, \check{X}) \end{bmatrix} \begin{bmatrix} \sigma(\check{c}, \check{x}) \\ \sigma^*(\check{X}) \end{bmatrix} \right) \\ &+ \alpha_h(\check{X}) \cdot \sigma(\check{c}, \check{x})' W_x(\check{x}, \check{X}) + \alpha_h(\check{X}) \cdot \sigma^*(\check{X})' W_X(\check{x}, \check{X}) \end{aligned}$$

Then  $\check{z}W(\check{x}, \check{X})$  the value function for the stochastic control problem:

$$\check{z}W(\check{x}, \check{X}) = E \int_0^\infty \exp(-\delta t) z_t U(c_t, x_t) dt$$

and evolution:

$$\begin{aligned} dx_t &= \mu(c_t, x_t)dt + \sigma(c_t, x_t)dB_t \\ dz_t &= z_t \alpha_h(X_t)dB_t \\ dX_t &= \mu^*(X_t)dt + \sigma^*(X_t)dB_t \end{aligned}$$

where  $z_0 = \check{z}$ ,  $x_0 = \check{x}$  and  $X_0 = \check{X}$ . To interpret the nonnegative  $z$  as inducing a change in probability, we initialize  $z_0$  at unity.

Also,  $W(\check{x}, \check{X}, \theta)$  is the value function for a control problem with discounted objective:

$$W(\check{x}, \check{X}) = \max_{c \in C} E \int_0^\infty \exp(-\delta t) U(c_t, x_t) dt$$

and evolution:

$$\begin{aligned} dx_t &= \mu(c_t, x_t)dt + \sigma(c_t, x_t) \left[ \alpha_h(X_t)dt + d\tilde{B}_t \right] \\ dX_t &= \mu^*(X_t)dt + \sigma^*(X_t) \left[ \alpha_h(X_t)dt + d\tilde{B}_t \right]. \end{aligned}$$

This value function is constructed using a perturbed specification where a Brownian increment  $dB_t$  is replaced by an increment  $\alpha_h(X_t)dt + d\tilde{B}_t$  with a drift distortion that depends only on the uncontrollable state  $X$ . This perturbation is justified via the Girsanov Theorem, provided that we entertain a weak solution to the stochastic differential equation governing the state evolution equation.

## References

- Anderson, E., L. P. Hansen, and T. Sargent (2003, March). A quartet of semi-groups for model specification, robustness, prices of risk, and model detection. *Journal of the European Economic Association* 1, 68–123.
- Araujo, A. and A. Sandroni (1999). On the convergence to homogeneous expectations when markets are complete. *Econometrica* 67(3), 663–672.
- Başar, T. and P. Bernhard (1995).  *$H_\infty$ -Optimal Control and Related Minimax Design Problems*. Boston: Birkhauser.
- Blackwell, D. and L. Dubins (1962). Merging of opinions with increasing information. *Annals of Mathematical Statistics* 38, 882–886.
- Blackwell, D. and M. Girshick (1954). *Theory of Games and Statistical Decisions*. New York: Wiley.
- Chamberlain, G. (2000). Econometric applications of maxmin expected utility theory. *Journal of Applied Econometrics* 15, 625–644.
- Duffie, D. and L. G. Epstein (1992). Stochastic differential utility. *Econometrica* 60(2), 353–394.
- Dupuis, P. and R. S. Ellis (1997). *A Weak Convergence Approach to the Theory of Large Deviations*. Wiley Series in Probability and Statistics. New York: John Wiley and Sons.
- Ekeland, I. and T. Turnbull (1983). *Infinite-Dimensional Optimization and Convexity*. Chicago Lectures in Mathematics. Chicago: The University of Chicago Press.
- Elliott, R. J. (1982). *Stochastic Calculus and Applications*. New York: Springer-Verlag.
- Epstein, L. and M. Schneider (2004). Recursive multiple-priors. *Journal of Economic Theory* 113(1), 1 – 31.
- Epstein, L. and S. Zin (1989). Substitution, risk aversion and the temporal behavior of consumption and asset returns: A theoretical framework. *Econometrica* 57, 937–969.
- Fan, K. (1952). Fixed-point and minimax theorems in locally convex topological linear spaces. *Proceedings of the National Academy of Sciences* 38, 121–126.
- Fan, K. (1953). Minimax theorems. *Proceedings of the National Academy of Sciences* 39, 42–47.
- Fleming, W. H. and H. M. Soner (1993). *Controlled Markov Processes and Viscosity Solutions*. Applications of Mathematics. New York: Springer-Verlag.
- Fleming, W. H. and P. E. Souganidis (1989). On the existence of value functions of two-player, zero sum stochastic differential games. *Indiana University Mathematics Journal* 38, 293 – 314.

- Föllmer, H. (1985). An entropy approach to the time reversal of diffusion processes. In M. Metivier and E. Pardoux (Eds.), *Stochastic Differential Systems*, Volume 69 of *Lecture Notes in Control and Information Sciences*, pp. 156–163. Springer-Verlag.
- Gilboa, I. and D. Schmeidler (1989). Maxmin expected utility with non-unique prior. *Journal of Mathematical Economics* 18, 141–153.
- Hansen, L. P. and T. Sargent (1995, May). Discounted linear exponential quadratic gaussian control. *IEEE Transactions on Automatic Control* 40(5), 968–971.
- Hansen, L. P., T. Sargent, and T. Tallarini (1999). Robust permanent income and pricing. *Review of Economic Studies* 66, 873–907.
- Hansen, L. P. and T. J. Sargent (2003, April). Robust control of forward-looking models. *Journal of Monetary Economics* 50(3), 581 – 604.
- Hansen, L. P. and T. J. Sargent (2005a). Robust estimation and control under commitment. unpublished.
- Hansen, L. P. and T. J. Sargent (2005b). Robust estimation and control without commitment. unpublished.
- Hansen, L. P. and T. J. Sargent (2006). *Robustness*. Princeton, New Jersey: Princeton University Press.
- Jackson, M. O., E. Kalai, and R. Smorodinsky (1999, July). Bayesian representation of stochastic processes under learning: De Finetti revisited. *Econometrica* 67(4), 875–893.
- Jacobson, D. H. (1973). Optimal stochastic linear systems with exponential performance criteria and their relation to deterministic differential games. *IEEE Transactions on Automatic Control* 18, 124–131.
- James, M. R. (1992). Asymptotic analysis of nonlinear stochastic risk sensitive control and differential games. *Mathematics of Control, Signals, and Systems* 5, 401–417.
- Kabanov, J. M., R. S. Lipcer, and A. N. Sirjaev (1979). Absolute continuity and singularity of locally absolutely continuous probability distributions. I. *Math. USSR Sbornik* 35, 631 – 696.
- Kalai, E. and E. Lerner (1993). Rational learning leads to Nash equilibrium. *Econometrica* 61(5), 1019–1045.
- Karatzas, I. and S. E. Shreve (1991). *Brownian Motion and Stochastic Calculus* (Second ed.). New York: Springer-Verlag.
- Kunita, H. (1969). Absolute continuity of Markov processes and generators. *Nagoya Mathematical Journal* 36, 1–26.
- Liptser, R. S. and A. N. Shiryaev (2000). *Statistics of Random Processes* (Second ed.), Volume I. General Theory of Applications of Mathematics. Berlin: Springer.
- Luenberger, D. G. (1969). *Optimization by Vector Space Methods*. New York: Wiley.

- Maccheroni, F., M. Marinacci, and A. Rustichini (2004). Variational representation of preferences under ambiguity. unpublished.
- Onatski, A. and N. Williams (2003). Modeling model uncertainty. *Journal of the European Economic Association* 1, 1087–1122.
- Petersen, I. R., M. R. James, and P. Dupuis (2000). Minimax optimal control of stochastic uncertain systems with relative entropy constraints. *IEEE Transactions on Automatic Control* 45, 398–412.
- Revuz, D. and M. Yor (1994). *Continuous Martingales and Brownian Motion* (Second ed.). New York: Springer-Verlag.
- Sandroni, A. (2000). Do markets favor agents able to make accurate predictions? *Econometrica* 68(6), 1303–1341.
- Schroder, M. and C. Skiadas (1999). Optimal consumption and portfolio selection with stochastic differential utility. *Journal of Economic Theory* 89(1), 68–126.
- Skiadas, C. (2001). Robust control and recursive utility. Department of Finance, Kellogg School of Management, Northwestern University.
- Wang, T. (2001). Two classes of multi-prior preferences. unpublished.
- Whittle, P. (1981). Risk-sensitive linear/quadratic/gaussian control. *Advances in Applied Probability* 13, 764–777.